

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

BA

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/00	A2	(11) International Publication Number: WO 00/06137 (43) International Publication Date: 10 February 2000 (10.02.00)
(21) International Application Number: PCT/US99/17267 (22) International Filing Date: 30 July 1999 (30.07.99) (30) Priority Data: 09/126,185 30 July 1998 (30.07.98) US (71) Applicant: ABBOTT LABORATORIES [US/US]; CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-6050 (US). (72) Inventors: COGHLAN, Michael, J.; 307 Fairfax Lane, Grayslake, IL 60030 (US). LULY, Jay, R.; 24 Damien Road, Wellesley, MA 02181 (US). SCHKERYANTZ, Jeffrey, M.; 526 English Lane, Winthrop Harbor, IL 60096 (US). WANG, Alan, X.; 1012 Talbot Lane, Grayslake, IL 60030 (US). (74) Agents: MILLER, Robert, A. et al.; Abbott Laboratories, CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US).		(81) Designated States: CA, JP, MX, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: GLUCOCORTICOID-SELECTIVE AGENTS <div style="text-align: center;"> <p style="margin-top: 10px;">(I)</p> </div> (57) Abstract <p>Compounds having Formula (I) are useful for modulating the glucocorticoid receptor in a mammal. Also disclosed are pharmaceutical compositions comprising compounds of Formula (I) and methods of treating immune, autoimmune, inflammatory, adrenal imbalance, cognitive and behavioral diseases in a mammal.</p> <div style="text-align: center; margin-top: 100px;"> <p>BEST AVAILABLE COPY</p> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

GLUCOCORTICOID-SELECTIVE AGENTS

Technical Field

The present invention relates to compounds which are selective for glucocorticoid receptors, pharmaceutical compositions comprising these compounds, and to methods of treating immune, autoimmune, inflammatory, adrenal imbalance, cognitive, and behavioral diseases.

5

Background of The Invention

Intracellular receptors (IR's) are a class of structurally related proteins involved in the regulation of gene expression. The steroid hormone receptors are a subset of this superfamily whose natural ligands are typically comprised of endogenous steroids such as estradiol, progesterone, and cortisol. Man-made ligands to these receptors play an important role in human health and, of these receptors, the glucocorticoid receptor (GR) has an essential role in regulating human physiology and immune response. Steroids which interact with GR have been shown to be potent antiinflammatory agents. Despite this benefit, steroidal GR ligands are not selective. Side effects associated with chronic dosing are believed to be the result of cross-reactivity with other steroid receptors such as estrogen, progesterone, androgen, and mineralocorticoid receptors which have homologous ligand binding domains.

A ligand which is selective for GR over other IRs could modulate (i.e. repress, agonize, partially agonize, or antagonize) and thus can be used to influence the basic, life-sustaining systems of the body including carbohydrate, protein and lipid metabolism and the functions of the cardiovascular, kidney, central nervous, immune, skeletal muscle, and other organ and tissue systems. In this regard, GR modulators have proven useful in the treatment of inflammation, tissue rejection, auto-immunity, malignancies such as leukemias and lymphomas, Cushing's syndrome, acute adrenal insufficiency, congenital adrenal hyperplasia, rheumatic fever, polyarteritis nodosa, granulomatous polyarteritis, inhibition of myeloid cell lines, immune proliferation/apoptosis, HPA axis suppression and regulation, hypercortisolemia, modulation of the Th1/Th2 cytokine balance, chronic kidney disease, stroke and spinal cord injury, hypercalcemia, hyperglycemia, acute adrenal insufficiency, chronic primary adrenal insufficiency, secondary adrenal insufficiency, congenital adrenal hyperplasia, cerebral edema, thrombocytopenia, and Little's syndrome.

GR modulators are especially useful in disease states involving systemic inflammation such as inflammatory bowel disease, systemic lupus erythematosus, polyarthritis nodosa, Wegener's granulomatosis, giant cell arteritis, rheumatoid arthritis, osteoarthritis, hay fever,

allergic rhinitis, urticaria, angioneurotic edema, chronic obstructive pulmonary disease, asthma, tendonitis, bursitis, Crohn's disease, ulcerative colitis, autoimmune chronic active hepatitis, organ transplantation, hepatitis, and cirrhosis. GR active compounds have also been used as immunostimulants, repressors, and wound healing and tissue repair agents.

5 GR modulators have also found use in a variety of topical diseases such as inflammatory scalp alopecia, panniculitis, psoriasis, discoid lupus erythematosus, inflamed cysts, atopic dermatitis, pyoderma gangrenosum, pemphigus vulgaris, bullous pemphigoid, systemic lupus erythematosus, dermatomyositis, herpes gestationis, eosinophilic fasciitis, relapsing polychondritis, inflammatory vasculitis, sarcoidosis, Sweet's disease, type 1 reactive leprosy, 10 capillary hemangiomas, contact dermatitis, atopic dermatitis, lichen planus, exfoliative dermatitis, erythema nodosum, acne, hirsutism, toxic epidermal necrolysis, erythema multiform, and cutaneous T-cell lymphoma.

Selective antagonists of the glucocorticoid receptor have been unsuccessfully pursued for decades. These agents would potentially find application in several disease states associated with 15 Human Immunodeficiency Virus (HIV), cell apoptosis, and cancer including, but not limited to, Kaposi's sarcoma, immune system activation and modulation, desensitization of inflammatory responses, IL-1 expression, anti-retroviral therapy, natural killer cell development, lymphocytic leukemia, and treatment of retinitis pigmentosa. Cognitive and behavioral processes are also susceptible to glucocorticoid therapy where antagonists would potentially be useful in the 20 treatment of processes such as cognitive performance, memory and learning enhancement, depression, addiction, mood disorders, chronic fatigue syndrome, schizophrenia, stroke, sleep disorders, and anxiety.

The prior art discloses a variety of triarylmethanes including triphenylmethanes useful as dyes or pigments. We have unexpectedly discovered a series of triphenylmethane compounds 25 which selectively modulate the glucocorticoid receptor in relation to the progesterone receptor, mineralocorticoid receptor, androgen receptor, and estrogen receptor-alpha. In addition, we provide a series of novel compounds which are selective for GR. Importantly, it has not been known previously that triphenylmethane compounds of this invention are useful as selective glucocorticoid receptor modulators.

30 Examples of the prior art include:

Aoyama et al. (European Patent application 85301391.0, published Sep. 04, 1985) discloses triarylmethane compounds which include triphenylmethane compounds that function as pigments during the process of determining the reduced form of nicotinamide adenine dinucleotide (phosphate);

Aoyama et al. (European Patent application 85302562.5, published Oct. 30, 1985) discloses triphenylmethane compounds that function as pigments during the process of determining compounds that contain a mercapto group;


Nieto et al. (Biochemistry International, 1990, Vol. 21, No. 2, 305-311) discloses that phenolphthalein, a dye containing a triphenylmethane structure, or a phenolphthalein derivative interacts with the rat estrogen receptor;

Kinoshita et al. (U.S. Patent No. 5,112,867, issued May 12, 1992) discloses triphenylmethane compounds that are useful for treating osteoporosis; and

Brugnara et al. (International Patent application 97/34589, published Sep. 25, 1997) discloses triphenylmethane compounds that have utility for inhibiting or treating sickle cell diseases and cell proliferation diseases in mammals.

Summary of The Invention

In the principle embodiment of the present invention is disclosed a method of selectively modulating the glucocorticoid receptor in a mammal comprising administering an effective amount of a compound of Formula I



I,

or a pharmaceutically acceptable salt or prodrug thereof, where

R^1 is selected from

- (1) hydrogen and
- (2) -OH;

L^1 is selected from

- (1) a covalent bond,
- (2) -O-,
- (3) -S(O)_t- where t is an integer from 0 to 2, and
- (4) -NR⁹- where R⁹ is selected from

- (a) hydrogen and
- (b) alkyl of one to four carbons;

R^2 and R^3 are independently selected from

- 5 (1) hydrogen,
- (2) an amino-protecting group,
- (3) alkyl of one to six carbons, and
- (4) alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from
- 10 (a) phenyl and
- (b) $-OR^{10}$ where R^{10} is selected from
 - (i) hydrogen,
 - (ii) alkyl of one to six carbons,
 - (iii) a hydroxy-protecting group, and
 - 15 (iv) $-C(O)R^{11}$ where R^{11} is selected from
 - alkyl of one to six carbons,
 - phenyl, and
 - phenyl substituted with 1, 2, or 3 substituents selected from
 - 20 $-NO_2$,
 - alkyl of one to six carbons, and
 - halogen, or

R^2 and R^3 together with the nitrogen to which they are attached form a 4 to 8
25 membered ring selected from the group consisting of heterocycle;

R^4 and R^5 are independently selected from

- (1) hydrogen,
- (2) halogen,
- 30 (3) $-NR^{12}R^{13}$ where R^{12} and R^{13} are independently selected from
 - (a) hydrogen,
 - (b) an amino-protecting group,
 - (c) alkyl of one to six carbons, and
 - (d) alkyl of one to six carbons substituted with 1, 2, or 3 substituents

- independently selected from
- (i) -OR¹⁰ and
 - (ii) phenyl, or
- R¹² and R¹³ together with the nitrogen to which they are attached form a
- 5 4 to 8 membered ring selected from the group consisting of heterocycle,
- (4) alkyl of one to six carbons, and
 - (5) alkyl of one to six carbons substituted with 1, 2, or 3 substituents
- independently selected from
- (a) halogen,
 - 10 (b) -OR¹⁰,
 - (c) -CN, and
 - (d) -CO₂R¹⁴ where R¹⁴ is selected from
- (i) hydrogen,
 - (ii) alkyl of one to six carbons, and
 - 15 (iii) alkyl of one to six carbons substituted with 1, 2, or 3
- substituents independently selected from
- phenyl and
- phenyl substituted with 1, 2, or 3 substituents
- independently selected from
- 20 -NO₂,
- alkyl of one to six carbons, and
- halogen, and
- (e) -NR¹⁵R¹⁶ where R¹⁵ and R¹⁶ are independently selected from
- (i) hydrogen,
 - 25 (ii) an amino-protecting group,
 - (iii) alkyl of one to six carbons, and
 - (iv) alkyl of one to six carbons substituted with 1, 2, or 3
- substituents independently selected from
- phenyl and
- 30 phenyl substituted with 1, 2, or 3 substituents
- independently selected from
- NO₂,
- alkyl of one to six carbons, and
- halogen;

R^6 , R^7 , and R^8 are independently selected from

- (1) hydrogen,
- (2) halogen,
- 5 (3) alkyl of one to six carbons,
- (4) alkyl of one to six carbons substituted with 1, 2, or 3 substituents
independently selected from
 - (a) halogen,
 - (b) $-OR^{10}$,
 - 10 (c) $-CN$,
 - (d) $-CO_2R^{14}$, and
 - (e) $-NR^{15}R^{16}$,
- (5) perfluoroalkyl of one to six carbons,
- (6) $-NR^{17}R^{18}$ where R^{17} and R^{18} are independently selected from
 - 15 (a) hydrogen,
 - (b) an amino-protecting group,
 - (c) alkyl of one to six carbons,
 - (d) alkyl of one to six carbons substituted with 1, 2, or 3 substituents
independently selected from
 - 20 (i) phenyl,
 - (ii) heterocycle, and
 - (iii) $-OR^{10}$,
 - (e) $-C(O)R^{19}$ where R^{19} is selected from
 - (i) alkyl of one to six carbons,
 - 25 (ii) alkyl of one to six carbons substituted with 1, 2, or 3
substituents independently selected from
heterocycle,
phenyl, and
phenyl substituted with 1, 2, or 3 substituents
30 independently selected from
alkyl of one to six carbons,
halogen,
 $-NO_2$,
 $-CF_3$,

- 5
10
15
20
25
30
- CN,
 - C(O)R²⁰ where R²⁰ is selected from
 - hydrogen,
 - alkyl of one to six carbons,
 - NR²¹R²² where R²¹ and R²² are independently selected from
 - hydrogen,
 - an amino-protecting group, and
 - alkyl of one to six carbons, and
 - OR²³ where R²³ is selected from
 - alkyl of one to six carbons and
 - alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from the group consisting of halogen,
 - (iii) cycloalkyl of three to six carbons,
 - (iv) cycloalkyl of three to six carbons substituted with 1, 2, or 3 substituents independently selected from
 - heterocycle,
 - phenyl,
 - phenyl substituted with 1, 2, or 3 substituents independently selected from
 - NO₂,
 - alkyl of one to six carbons, and
 - halogen,
 - halogen,
 - CN,
 - CO₂R¹⁴,
 - alkyl of one to six carbons, and
 - alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from the group consisting of halogen,
 - (v) alkenyl of two to six carbons,
 - (vi) alkenyl of two to six carbons substituted with 1 or 2


- substituents independently selected from
heterocycle,
phenyl, and
phenyl substituted with 1, 2, or 3 substituents
independently selected from
alkyl of one to six carbons,
halogen,
-NO₂,
-CF₃,
-CN, and
-CO₂R¹⁴,
- (vii) phenyl,
(viii) phenyl substituted with 1, 2, or 3 substituents independently
selected from
halogen,
alkyl of one to six carbons,
-NO₂,
-CF₃,
-CN, and
-CO₂R¹⁴, and
- (ix) -OR¹¹, and
(f) -SO₂R²⁴ where R²⁴ is selected from
(i) alkyl of one to six carbons,
(ii) phenyl, and
(iii) phenyl substituted with 1, 2, or 3 substituents independently
selected from
alkyl of one to six carbons and
-NO₂,
- (7) -OR²⁵ where R²⁵ is selected from
(a) perfluoroalkyl of one to six carbons,
(b) alkyl of one to six carbons,
(c) alkyl of one to six carbons substituted with 1, 2, or 3 substituents
independently selected from
(i) halogen and

- (ii) phenyl,
- (d) a hydroxy-protecting group, and
- (e) $-C(O)R^{14}$,
- (8) $-CN$,
- 5 (9) $-C(O)R^{19}$,
- (10) $-CO_2R^{14}$,
- (11) $-C(O)R^{20}$,
- (12) $-SO_2NR^{26}R^{27}$ where R^{26} and R^{27} are independently selected from,
 - (a) alkyl of one to six carbons,
 - 10 (b) phenyl, and
 - (c) phenyl substituted with 1, 2, or 3 substituents independently selected from
 - (i) alkyl of one to six carbons,
 - (ii) halogen, and
 - 15 (iii) $-NO_2$,
- (13) $-S(O)_tR^{28}$ where t is defined previously and R^{28} is selected from
 - (a) hydrogen,
 - (b) perfluoroalkyl of one to six carbons,
 - (c) alkyl of one to six carbons,
 - 20 (d) alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from
 - (i) phenyl and
 - (ii) phenyl substituted with 1, 2, or 3 substituents independently selected from
 - 25 alkyl of one to six carbons,
 - halogen, and
 - $-NO_2$,
 - (e) phenyl, and
 - (f) phenyl substituted with 1, 2, or 3 substituents independently selected from
 - 30 (i) alkyl of one to six carbons,
 - (ii) halogen, and
 - (iii) $-NO_2$,
- (14) $-NO_2$,

(15) $-N=CHR^{29}$ where R^{29} is selected from

- (a) phenyl,
- (b) aryl, and
- (c) heterocycle, and

5 (16)



where X is selected from $-\text{CH}_2-$, $-\text{CH}_2\text{O}-$ and $-\text{O}-$, and Y is selected from $-\text{C}(\text{O})-$ and $-(\text{C}(\text{R}'')_2)_v-$, where R'' is hydrogen or alkyl of one to four carbons, and v is an integer from 1 to 3.

10 In still another embodiment of the present invention is disclosed a method of treating inflammation and immune, autoimmune and inflammatory diseases in a mammal comprising administering an effective amount of a compound of Formula I.

In still another embodiment of the present invention is disclosed a method of treating adrenal imbalance in a mammal comprising administering an effective amount of a compound of Formula I.

15 In still another embodiment of the present invention is disclosed a method of treating cognitive and behavioral processes susceptible to glucocorticoid therapy where antagonists would be useful in the treatment of processes such as cognitive performance, memory and learning enhancement, depression, addiction, mood disorders, chronic fatigue syndrome, schizophrenia, stroke, sleep disorders, and anxiety in a mammal comprising administering an effective amount of a compound of Formula I.

20 In still another embodiment of the present invention is disclosed pharmaceutical compositions containing compounds of Formula I.

Compounds of this invention include, but are not limited to,

- 4',4" bis(dimethylamino)-2-chloro-5-nitrotriphenylmethane,
- 25 4',4" bis(dimethylamino)-4-chloro-3-nitrotriphenylmethane,
- 4',4" bis(dimethylamino)-5-acetamido-2-chlorotriphenylmethane,
- 4',4" bis(dimethylamino)-4-nitrotriphenylmethane ,
- 4',4" bis(dimethylamino)-4-chlorotriphenylmethane,
- 4',4" bis(dimethylamino)-3-chlorotriphenylmethane,
- 30 4',4" bis(dimethylamino)-2-chlorotriphenylmethane,
- 4',4" bis(dimethylamino)-2-methoxytriphenylmethane,
- 4',4" bis(dimethylamino)-3-nitrotriphenylmethane,

- 4',4'' bis(dimethylamino)-2-trifluoromethyltriphenylmethane,
4',4'' bis(dimethylamino)triphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-6-nitrotriphenylmethane,
4',4'' bis(N-piperidinyl)-2-chloro-5-nitrotriphenylmethane,
5 4',4'' bis(dimethylamino)-2-bromotriphenylmethane,
4',4'' bis(dimethylamino)-2-methyltriphenylmethane,
4',4'' bis(dimethylamino)-2,3,5-trichlorotriphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-5-trifluoromethyltriphenylmethane,
4',4'' bis(dimethylamino)-2,4-dichlorotriphenylmethane,
10 4',4'' bis(dimethylamino)-2-chloro-4,5-methylenedioxytriphenylmethane,
4',4'' bis(dimethylamino)-2,6-dichlorotriphenylmethane,
4',4'' bis(dimethylamino)-2,3-dichlorotriphenylmethane,
4',4'' bis(N-morpholinyl)-2-chloro-5-nitrotriphenylmethane,
4',4'' bis(methylamino)-2-chloro-5-nitrotriphenylmethane,
15 4',4'' bis(dimethylamino)-2,5-dichlorotriphenylmethane,
4',4'' bis(dimethylamino)-2-fluorotriphenylmethane,
4',4'' bis(dimethylamino)-2-iodotriphenylmethane,
4',4'' bis(methylamino)-2,5-dichlorotriphenylmethane,
4',4'' bis(N-morpholinyl)-2,3,5-trichlorotriphenylmethane,
20 4',4'' bis(N-pyrrolidinyl)-2-chloro-5-nitrotriphenylmethane,
4',4'' bis(di-n-butylamino)-2-chloro-5-nitrotriphenylmethane,
4',4'' bis(N-(2-acetoxyethyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane,
4',4'' bis(N-(2-hydroxyethyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-5-iodotriphenylmethane,
25 4',4'' bis(dimethylamino)-5-bromo-2-chlorotriphenylmethane,
4',4'' bis(N-(t-butoxycarbonyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane,
4',4'' bis(N-benzylamino)-2-chloro-5-nitrotriphenylmethane,
4',4'' bis(N-benzylamino)-2,5-dichlorotriphenylmethane,
4',4'' bis(dimethylamino)-4-methoxytriphenylmethane,
30 4'-dimethylamino-4''-methylamino-2-chloro-5-nitrotriphenylmethane,
4'-dimethylamino-4''-(N-morpholinyl)-2-chloro-5-nitrotriphenylmethane,
4'-dimethylamino-4''-(N-pyrrolidinyl)-2-chloro-5-nitrotriphenylmethane,
4'-dimethylamino-4''-(N-methyl-N-(2-hydroxyethyl)amino)-2-chloro-5-nitrotriphenylmethane,

- 4'-dimethylamino-4''-(di-n-butylamino)-2-chloro-5-nitrotriphenylmethane,
 3'-methyl-4',4''-bis(dimethylamino)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-piperidinyl)-2-chloro-5-nitrotriphenylmethane,
 4'-methylamino-4''-(t-butoxycarbonylamino)-2-chloro-5-nitrotriphenylmethane,
 5 4'-methylamino-4''-(N-(2-acetoxyethyl)-N-methylamino)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-methyl-N-(2-benzoyloxyethyl)amino)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-methyl-N-(2-acetoxyethyl)amino)-2-chloro-5-nitrotriphenylmethane,
 10 phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] ether,
 4-chlorophenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] ether,
 phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] thioether,
 phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] amine,
 15 4'-dimethylamino-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-2-chlorotriphenylmethanol,
 4'-dimethylamino-2-chlorotriphenylmethane,
 4',4'' bis(dimethylamino)-5-amino-2-chlorotriphenylmethane,
 4',4'' bis(dimethylamino)-2-chloro-5-(4-nitrobenzamido)triphenylmethane,
 20 4',4'' bis(dimethylamino)-2-chloro-5-(4-nitrocinnamido)triphenylmethane,
 4',4'' bis(dimethylamino)-2-chloro-5-(cyclopropylcarbamido)triphenylmethane,
 4',4'' bis(dimethylamino)-2-chloro-5-(dimethylsulphonimido)triphenylmethane,
 4',4'' bis(dimethylamino)-2-chloro-5-(methoxycarbonylamino)triphenylmethane,
 4',4'' bis(dimethylamino)-2-chloro-5-(2-furanylmethylimino)triphenylmethane, and
 25 4',4'' bis(dimethylamino)-2-chloro-5-(2-furanylmethylamino)triphenylmethane.

Detailed Description of The Invention

Definition of Terms

- The term "alkenyl of two to six carbons" refers to a straight or branched chain
 30 hydrocarbon radical containing from two-to-six carbon atoms and also containing at least one
 carbon-carbon double bond. Representative examples of "alkenyl of two to six carbons"
 include but are not limited to groups such as ethenyl, propenyl, isobutenyl, 1-butenyl, 1,3-
 butadienyl, 2-pentenyl, 2-hexenyl, 1,5-hexadienyl and the like.

The term "alkyl of one to four carbons" refers to a straight or branched chain hydrocarbon radical containing from one-to-four carbon atoms. Representative examples of "alkyl of one to four carbons" include but are not limited to groups such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, tert-butyl and the like. The term "alkyl of one to six carbons" refers to a straight or branched chain hydrocarbon radical containing from one-to-six carbon atoms. Representative examples of "alkyl of one to six carbons" include but are not limited to groups such as all of the previous examples as well as n-pentyl, isopentyl, neopentyl, n-hexyl and the like.

The term "amino" refers to -NH_2 .

The term "amino-protecting group" refers to groups intended to protect an amino group against undesirable reactions during synthetic procedures. Commonly used N-protecting groups are disclosed in Greene, T. W., & Wuts, P. G. M. (1991). Protective Groups In Organic Synthesis (2nd ed.). New York: John Wiley & Sons. Preferred N-protecting groups are formyl, acetyl, benzoyl, pivaloyl, t-butylacetyl, phenylsulfonyl, benzyl, t-butyloxycarbonyl (Boc), and benzyloxycarbonyl (Cbz).

The term "aryl" as used herein refers to a carbocyclic ring system having 6-10 ring atoms and one or two aromatic rings. Representative examples of aryl groups include groups such as, for example, phenyl, naphthyl, tetrahydronaphthyl, indanyl, indenyl and the like. The aryl groups of this invention can be optionally substituted.


The term "cycloalkyl of three to six carbons" refers to a saturated cyclic hydrocarbon radical containing from three-to-six carbon atoms. Representative examples of "cycloalkyl of three to six carbons" include but are not limited to groups such as, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the like.

The term "halogen" refers to F, Cl, Br, or I.


The term "heterocycle" represents a represents a 4-, 5-, 6-, 7-, or 8-membered ring containing one, two or three heteroatoms independently selected from the group consisting of nitrogen, oxygen and sulfur. The 4- and 5-membered rings have zero to two double bonds, the 6- and 7-membered rings have zero to three double bonds and the 8-membered rings have zero to four double bonds. The term "heterocycle" also includes bicyclic, tricyclic and tetracyclic groups in which any of the above heterocyclic rings is fused to one or two rings independently selected from an aryl ring, a cyclohexane ring, a cyclohexene ring, a cyclopentane ring, a cyclopentene ring or another monocyclic heterocyclic ring. Heterocycles include acridinyl, benzimidazolyl, benzofuryl, benzothiazolyl, benzothienyl, benzoxazolyl, biotinyl, cinnolinyl, dihydrofuryl, dihydroindolyl, dihydropyranyl, dihydrothienyl, dithiazolyl, furyl,

homopiperidinyl, imidazolidinyl, imidazolynyl, imidazolyl, indolyl, isoquinolyl, isothiazolidinyl, isothiazolyl, isoxazolidinyl, isoxazolyl, morpholinyl, oxadiazolyl, oxazolidinyl, oxazolyl, piperazinyl, piperidinyl, pyranyl, pyrazolidinyl, pyrazinyl, pyrazolyl, pyrazolynyl, pyridazinyl, pyridyl, pyrimidinyl, pyrimidyl, pyrrolidinyl, pyrrolinyl, pyrrolyl, quinolynyl, quinoxaloyl, tetrahydrofuryl, tetrahydroisoquinolyl, tetrahydroquinolyl, tetrazolyl, thiadiazolyl, thiazolidinyl, thiazolyl, thienyl, thiomorpholinyl, triazolyl, and the like.

Heterocyclics also include bridged bicyclic groups where a monocyclic heterocyclic group is bridged by an alkylene group such as



Heterocyclics also include compounds of the formula



where X is selected from $-\text{CH}_2-$, $-\text{CH}_2\text{O}-$ and $-\text{O}-$, and Y is selected from $-\text{C}(\text{O})-$ and $-(\text{C}(\text{R}'')_2)_v-$, where R'' is hydrogen or alkyl of one to four carbons, and v is 1-3. These heterocycles include 1,3-benzodioxolyl, 1,4-benzodioxanyl, and the like.

15 The term "hydroxy-protecting group" refers to a substituent which protects hydroxyl groups against undesirable reactions during synthetic procedures. Examples of hydroxy-protecting groups include, but are not limited to, ethers, for example, methyl, ethyl, t-butyl, benzyl and allyl; substituted methyl ethers, for example, methoxymethyl, benzyloxymethyl, 2-methoxyethoxymethyl, 2-(trimethylsilyl)-ethoxymethyl, and triphenylmethyl; substituted ethyl
20 ethers, for example, 2,2,2-trichloroethyl and t-butyl; tetrahydropyranyl ethers; silyl ethers, for example, trimethylsilyl, t-butyldimethylsilyl and t-butyldiphenylsilyl; esters, for example, formate, acetate, trifluoroacetate, pivalate, benzoate, and adamantate; carbonates, for example, methyl, ethyl, isobutyl, t-butyl, vinyl, allyl, and benzyl; sulfonates, for example, methanesulfonate, benzy sulfonate and p-toluenesulfonate. Commonly used hydroxy-
25 protecting groups are disclosed in Greene, T. W., & Wuts, P. G. M. (1991). Protective Groups In Organic Synthesis (2nd ed.). New York: John Wiley & Sons.

The term "Lewis acid" refers to any chemical species which has a vacant orbital and therefore acts as an electron pair acceptor. Representative examples of a "Lewis acid" include but are not limited to boron trifluoride, aluminum trichloride, titanium tetrachloride, and stannic
30 tetrachloride.

The term "perfluoroalkyl of one to six carbons" refers to an alkyl group containing one-to-six carbon atoms where all the hydrogens have been substituted with fluorides.

The term "pharmaceutically acceptable prodrugs" represents those prodrugs of the compounds of the present invention which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals with undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention.

The term "prodrug" represents compounds which are rapidly transformed in vivo to the parent compound of the above formula, for example, by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, Vol. 14 of the A.C.S. Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference.

The term "pharmaceutically acceptable salt" represents those salts which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge, *et al.* describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, **1977**, 66:1 - 19. The salts can be prepared *in situ* during the final isolation and purification of the compounds of the invention, or separately by reacting the free base function with a suitable organic acid. Representative acid addition salts include acetate, adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphersulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptonate, glycerophosphate, hemisulfate, heptonate, hexanoate, hydrobromide, hydrochloride, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, toluenesulfonate, undecanoate, valerate salts, and the like. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like, as well as nontoxic ammonium, quaternary ammonium, and amine cations, including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like.

Compounds of the present invention can exist as stereoisomers where asymmetric or chiral centers are present. These compounds are designated by the symbols "R" or "S," depending on the configuration of substituents around the chiral carbon atom. The present invention contemplates various stereoisomers and mixtures thereof. Stereoisomers include enantiomers and diastereomers, and equal mixtures of enantiomers are designated (\pm). Individual stereoisomers of compounds of the present invention can be prepared synthetically from commercially available starting materials which contain asymmetric or chiral centers or by preparation of racemic mixtures followed by resolution well-known to those of ordinary skill in the art. These methods of resolution are exemplified by (1) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography and liberation of the optically pure product from the auxiliary or (2) direct separation of the mixture of enantiomers on chiral chromatographic columns.

15 Determination of Biological Activity

For glucocorticoid receptor (GR) cytosol binding assays, the procedure described in Anal. Biochem. 1970, 37, 244-252, hereby incorporated by reference, was used. Briefly, cytosol preparations of human glucocorticoid receptor- α [GR α] isoform and human progesterone receptor-A [PRA] isoform were obtained from Ligand Pharmaceuticals (San Diego, CA). Both receptor cDNAs were cloned into baculovirus expression vectors and expressed in insect SF21 cells. [3 H]-dexamethasone (Dex, specific activity 82-86 Ci/mmol) and [3 H]-progesterone (Prog, specific activity 97-102 Ci/mmol) were purchased from Amersham Life Sciences (Arlington Heights, IL). Glass fiber type C multiscreen MAFC NOB plates were from Millipore (Burlington, MA). Hydroxyapatite Bio-Gel HTP gel was obtained from Bio-Rad Laboratories (Hercules, CA). Tris(hydroxymethyl)aminomethane (Tris), ethylenediaminetetraacetic acid (EDTA), glycerol, dithiothreitol (DTT) and sodium molybdate were obtained from Sigma Chemicals (St. Louis, MO). Microscint-20 scintillation fluid was obtained from Packard Instrument (Meriden, CT).

30 Stock solutions (32 mM) of compounds were prepared in dimethylsulfoxide (DMSO), and 50X solutions of test compounds were prepared from the 32 mM solution with a 50:50 mixture of DMSO/ethanol. The 50X solution was then diluted with binding buffer that contained 10 mM Tris-HCl, 1.5 mM EDTA, 10% glycerol, 1 mM DTT, 20 mM sodium molybdate, pH 7.5 at 4 °C. 1% DMSO/ethanol was also present in the binding assay.

GRX and PRA binding reactions were performed in Millipore Multiscreen plates. For GR binding assays, [^3H]-Dex (~35,000 dpm (~0.9 nM)), GRX cytosol (~35 μg protein), test compounds and, binding buffer were mixed in a total volume of 200 μL and incubated at 4 $^{\circ}\text{C}$ overnight in a plate shaker. Specific binding was defined as the difference between binding of [^3H]Dex in the absence and in the presence of 1 μM unlabeled Dex.

For progesterone receptor cytosol (PR) binding assays, [^3H]Prog (~36,000 dpm (~0.8 nM)), PRA cytosol (~40 μg protein), test compounds and binding buffer were mixed in a total volume of 200 μL and incubated at 4 $^{\circ}\text{C}$ overnight in a plate shaker. Specific binding was defined as the difference between binding of [^3H]Prog in the absence and in the presence of 3 μM unlabeled Prog.

After an overnight incubation, 50 μL of hydroxyapatite (25 % weight/volume) slurry were added to each well and plates were incubated for 10 min at 4 $^{\circ}\text{C}$ in a plate shaker. Plates were suctioned with a Millipore vacuum manifold and each well was rinsed with 300 μL of ice-cold binding buffer. A 250 μL aliquot of Packard Microscint-20 was added to each well and the wells were shaken at room temperature for 20 minutes. The amount of radioactivity was determined with a Packard TopCount plate reader.

For the determination of GR and PR inhibition constants (K_i), the concentration of test compounds that inhibited 50% of specific binding (IC_{50}) was determined from a Hill analysis of the competitive binding experiments. The K_i of test compounds was determined using the Cheng-Prusoff equation $K_i = \text{IC}_{50} / (1 + [L^*] / K_L)$ where L^* is the concentration of radioligand and K_L is the dissociation constant of the radioligand determined from saturation analysis. For GRX, K_L was ~1.5 nM, and for PRA, K_L was ~4.5 nM.

For soluble mineralocorticoid receptor (MR) binding assays, [^3H]-aldosterone (75-85 Ci/mmol) was purchased from New England Nuclear (Boston, MA), aldosterone was purchased from Sigma Chemical Co. (St. Louis, MO), CHAPS and DTT were purchased from Boehringer Mannheim GmbH (W. Germany) and all other reagents were purchased from Sigma.

The full length human androgen receptor was derived from cDNA expressed in a baculovirus expression system. The methods concerning growth, purification, and assays of recombinant viruses followed the protocol outlined by Summers, M.D., Smith, G.E., A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures, Tex. Agric. Exp. Stn. [Bull], 1987, No. 155, hereby incorporated by reference. The recombinant plasmids were cotransfected into SF21 cells with wild type AcNPV DNA, and the recombinant viruses were plaque purified.

A receptor extract was prepared from the baculovirus system, and aliquots were stored at -80 °C until used. Typical protein concentrations for these extracts were between 10 and 20 mg/ml. Stock solutions of aldosterone or other competing compounds were prepared as either 5 mM ethanol or DMSO stock solutions and serial dilutions were carried out in 1:1 DMSO-ethanol. The assay buffer consisted of the following: 25 mM sodium phosphate, 10 mM potassium fluoride, 20 mM sodium molybdate, 10% glycerol, 2 mM DTT and 0.25 mM CHAPS, pH=7.3 at room temperature.

Receptor assays were performed with a final volume of 250 µL containing from 50-75 µg of extract protein, plus 3-4 nM [³H]-aldosterone and varying concentrations of competing ligand (0 to 10,000 nM). Assays were set up using a 96-well minitube system and incubations were carried out at 4 °C for 18 hours. Equilibrium under these conditions of buffer and temperature was achieved by 6-8 hours. Nonspecific binding was defined as that binding remaining in the presence of 1000 nM unlabeled aldosterone. At the end of the incubation period, 200 µL of 6.25% hydroxyapatite was added in wash buffer (binding buffer in the absence of DTT and CHAPS). Specific ligand binding to receptor was determined by a hydroxyapatite-binding assay according to the protocol outlined by Weeksler, W.R., Norman, A.W., An Hydroxylapatite Batch Assay for the Quantitation of 1α, 25-Dihydroxyvitamin D₃-Receptor Complexes, *Anal. Biochem.* 1979, 92, 314-323, hereby incorporated by reference. Hydroxyapatite absorbs the receptor-ligand complex, allowing for the separation of bound from free radiolabeled ligand. The mixture was vortexed, incubated for 10 minutes at 4 °C, centrifuged, and the supernatant was removed. The hydroxyapatite pellet was washed twice more with the wash buffer. The amount of receptor-ligand complex was determined by liquid scintillation counting of the hydroxyapatite pellet after the addition of 0.5 mM EcoScint A scintillation cocktail from National Diagnostics (Atlanta, GA.).

After correcting for nonspecific binding, IC₅₀ values were determined. The IC₅₀ value is defined as the concentration of competing ligand required to decrease specific binding by 50%. The IC₅₀ values were determined graphically from a log-logit plot of the data. K_i values for the analogs were calculated by application of the Cheng-Prusoff equation outlined by Cheng, Y.-C., Prusoff, W.F., Relationship Between the Inhibition Constant (K_i) and the Concentration of Inhibitor Which Causes 50% Inhibition (IC₅₀) of an Enzymatic Reaction, *Biochem. Pharmacol.* 1973, 22, 3099-3108, hereby incorporated by reference. Standards are included in each assay and resulting K_i values are determined by use of a modified Cheng-Prusoff equation and used to calculate K_i values for "unknown analogs" as outlined by DeBlasi, A., O'Reilly, K., Motulsky, H.J., Calculating Receptor Number from Binding Experiments

Using Same Compound As Radioligand and Competitor, TIPS 1989, 10, 227-229, hereby incorporated by reference.

For human androgen receptor (AR) cytosol binding assays, [³H]-dihydrotestosterone (DHT) (120-140 Ci/mmol) was purchased from Amersham Life Science (Arlington Heights, IL), DHT was purchased from Sigma Chemical Co. (St. Louis, MO), CHAPS and DTT were purchased from Boehringer Mannheim GmbH (W. Germany), and all other reagents were purchased from Sigma. The full length human androgen receptor (AR) was derived from cDNA expressed in a baculovirus expression system from Ligand Pharmaceuticals (San Diego, CA). The methods concerning growth, purification, and assays of recombinant viruses followed the protocol outlined by Summers, M.D., Smith, G.E., A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures, Tex. Agric. Exp. Stn. [Bull], 1987, No. 155, hereby incorporated by reference. The recombinant plasmids were cotransfected into SF21 cells with wild type AcNPV DNA, and the recombinant viruses were plaque purified.

A receptor extract was prepared from the baculovirus system, and aliquots were stored at -80 °C until used. Typical protein concentrations for these extracts were between 10 and 20 mg/ml. Stock solutions of DHT or other competing compounds were prepared as 5 mM stock solutions in either 100% ethanol or DMSO and serial dilutions were carried out in 1:1 DMSO-ethanol. The assay buffer consisted of the following: 25 mM sodium phosphate, 10 mM potassium fluoride, 10 mM sodium molybdate, 10% glycerol, 1.5 mM EDTA, 2 mM DTT, 2 mM CHAPS and 1 mM PMSF, pH=7.4 at room temperature.

Receptor assays were performed with a final volume of 250 µL containing from 50-75 µg of extract protein, plus 1-2 nM [³H]-DHT and varying concentrations of competing ligand (0-10-5 M). Assays were set up using a 96-well minitube system and incubations were carried out at 4 °C for 18 hours. Equilibrium under these conditions of buffer and temperature was achieved by 6-8 hours. Nonspecific binding was defined as that binding remaining in the presence of 1000 nM unlabeled DHT. At the end of the incubation period, 200 µL of 6.25% hydroxyapatite was added in wash buffer (binding buffer in the absence of DTT and PMSF). Specific ligand binding to receptor was determined by a hydroxyapatite-binding assay according to the protocol outlined by Weckslar, W.R., Norman, A.W., An Hydroxylapatite Batch Assay for the Quantitation of 1α, 25-Dihydroxyvitamin D₃-Receptor Complexes, Anal. Biochem. 1979, 92, 314-323, hereby incorporated by reference. Hydroxyapatite absorbs the receptor-ligand complex, allowing for the separation of bound from free radiolabeled ligand. The mixture was vortexed, incubated for 10 minutes at 4 °C, centrifuged, and the supernatant was removed. The hydroxyapatite pellet was washed twice more with the wash buffer. The amount

of receptor-ligand complex was determined by liquid scintillation counting of the hydroxyapatite pellet after the addition of 0.5 mM EcoScint A scintillation cocktail from National Diagnostics (Atlanta, GA.).

After correcting for nonspecific binding, IC_{50} values were determined. The IC_{50} value is defined as the concentration of competing ligand required to decrease specific binding by 50%. The IC_{50} values were determined graphically from a log-logit plot of the data. K_i values for the analogs were calculated by application of the Cheng-Prusoff equation outlined by Cheng, Y.-C., Prusoff, W.F., Relationship Between the Inhibition Constant (K_i) and the Concentration of Inhibitor Which Causes 50% Inhibition (IC_{50}) of an Enzymatic Reaction, Biochem. Pharmacol. 1973, 22, 3099-3108, hereby incorporated by reference. Standards are included in each assay and resulting K_i values are determined by use of a modified Cheng-Prusoff equation and used to calculate K_i values for "unknown analogs" outlined by DeBlasi, A., O'Reilly, K., Motulsky, H.J., Calculating Receptor Number from Binding Experiments Using Same Compound As Radioligand and Competitor, TIPS 1989, 10, 227-229, hereby incorporated by reference.

For soluble estrogen receptor-alpha ($ER-\alpha$) binding assays, [3H]- Estradiol (120-140 Ci/mmol) was purchased from New England Nuclear (Boston, MA), 17-beta estradiol was purchased from Sigma Chemical Co. (St. Louis, MO), CHAPS and DTT were purchased from Boehringer Mannheim GmbH (W. Germany) and all other reagents were purchased from Sigma.

The human estrogen receptor-alpha cDNA was cloned into a yeast vector termed pYhER α and used to transform wild type yeast strain BJ2168 following the protocol outlined by Pham, T.A., Hwung Y.P., Santiso-Mere, D., McDonnell D.P., O'Malley, B.W., Ligand-Dependent and Independent Function of the Transactivation Regions of the Human Estrogen Receptor in Yeast, Mol. Endocrinol. 1992, 6, 1043-1050, hereby incorporated by reference. The yeast was induced with copper for 16 hours after which the cells were harvested, washed and the receptor extract prepared in cold buffer via a Bead Beater (BioSpec Products, Bartlesville, OK). Aliquots of the receptor typically contained 5-10 mg/ml of total protein and were stored at -80 °C until used. Estradiol or other competing compounds were prepared as 5 mM stock solutions in either 100% ethanol or DMSO and serial dilutions were carried out in 1:1 DMSO-ethanol. The assay buffer consisted of the following: 300 mM potassium chloride, 10 mM Trizma Base, 2 mM DTT and 5 mM CHAPS, pH=7.5 at room temperature.

Receptor assays were performed in a 250 μ L final volume containing from 5-10 μ g of extract protein, plus 2-3 nM [3H]-estradiol and varying concentrations of competing ligand (0 to

- 10,000 nM). Assays were set up using a 96-well minitube system and incubations were carried out at 4 °C for 18 hours. Equilibrium under these conditions of buffer and temperature was achieved by 6-8 hours. Nonspecific binding was defined as that binding remaining in the presence of 1000 nM unlabeled estradiol. At the end of the incubation period, 200 µL of 6.25% hydroxyapatite was added in wash buffer (binding buffer in the absence of DTT but containing 1 mM CHAPS). Specific ligand binding to receptor was determined by a hydroxyapatite-binding assay according to the protocol outlined by Weckslar, W.R., Norman, A.W, An Hydroxylapatite Batch Assay for the Quantitation of 1α, 25-Dihydroxyvitamin D₃-Receptor Complexes, Anal. Biochem. 1979, 92, 314-323, hereby incorporated by reference.
- Hydroxyapatite absorbs the receptor-ligand complex, allowing for the separation of bound from free radiolabeled ligand. The mixture was vortexed, incubated for 10 minutes at 4 °C, centrifuged, and the supernatant was removed. The hydroxyapatite pellet was washed twice more with the wash buffer. The amount of receptor-ligand complex was determined by liquid scintillation counting of the hydroxyapatite pellet after the addition of 0.5 mM EcoScint A scintillation cocktail from National Diagnostics (Atlanta, GA.).

- After correcting for nonspecific binding, IC₅₀ values were determined. The IC₅₀ value is defined as the concentration of competing ligand required to decrease specific binding by 50%. The IC₅₀ values were determined graphically from a log-logit plot of the data. K_i values for the analogs were calculated by application of the Cheng-Prusoff equation, Cheng, Y.-C., Prusoff, W.F., Relationship Between the Inhibition Constant (K_i) and the Concentration of Inhibitor Which Causes 50% Inhibition (IC₅₀) of an Enzymatic Reaction., Biochem. Pharmacol. 1973, 22, 3099-3108, hereby incorporated by reference. Standards are included in each assay and resulting K_i values are determined by use of a modified Cheng-Prusoff equation and used to calculate K_i values for "unknown analogs", DeBlasi, A., O'Reilly, K., Motulsky, H.J., Calculating Receptor Number from Binding Experiments Using Same Compound As Radioligand and Competitor, TIPS 1989, 10, 227-229, hereby incorporated by reference.

- The K_i of test compounds for MR, AR, and ER-α were determined using the Cheng-Prusoff equation, $K_i \text{ analog} = IC_{50 \text{ analog}} / (1 + [L] / K_d(L))$ where [L] is the labeled ligand concentration, K_d(L) is the K_d of the labeled ligand, and IC₅₀ is the concentration of analog to displace 50% of labeled ligand (determined graphically by log-logit plot of binding curve).

The inhibitory potencies of compounds of this invention and their selectivity for GR, PR, MR, AR, and ER-α receptors are shown in Table 1.

Table 1

Example Number	Ki (nM)				
	GR	PR	MR	AR	ER- α
1	272	>10,000	2899	3731	-
2	4200	-	-	-	-
3	5154	-	-	-	-
4	4649	-	-	-	-
5	4649	-	-	-	-
6	2525	-	-	-	-
7	1992	-	-	-	-
8	4288	-	-	-	-
9	1385	-	-	-	-
10	2228	-	-	-	-
11	4649	-	-	-	-
12	3852	-	-	-	-
13	5002	-	-	-	-
14	1514	-	-	-	-
15	2961	-	-	-	-
16	479	-	-	-	-
17	4704	-	-	-	-
18	5002	-	-	-	-
19	1600	-	-	-	-
20	1930	-	-	-	-
21	1891	-	-	-	-
22	386				
23	60	624	>10,000	4258	>10,000
24	306	2591	2899	3731	>10,000
25	5272	-	-	-	-
26	734	-	-	-	-

27	71	318	>10,000	3243	>10,000
28	5272	-	-	-	-
29	4561	-	-	-	-
31	145	2681	3906	3521	>10,000
32	136	536	3906	3521	>10,000
33	2304	-	-	-	-
34	725	-	-	-	-
35	4561	-	-	-	-
36	38	2591	2899	3731	>10,000
37	421	-	-	-	-
38	4712	-	-	-	-
39	127	510	2899	3731	>10,000
40	252	-	-	-	-
41	4494	-	-	-	-
42	109	187	2899	2797	>10,000
43	4494	-	-	-	-
44	287	-	-	-	-
46	1161	-	-	-	-
48	852	-	-	-	-
49	135	241	3906	1039	>10,000
50	1627	-	-	-	-
51	167	1084	>10,000	>10,000	>10,000
52	733	698	3105	>10,000	>10,000
53	788	-	-	-	-
54	227	2591	2899	3731	>10,000
55	1685	-	-	-	-
56	1026	-	-	-	-
57	2029	-	-	-	-
58	4303	-	-	-	-

59	4595	-	-	-	-
60	4651	-	-	-	-
61	4595	-	-	-	-
62	4595	-	-	-	-
63	4595	-	-	-	-
64	5272	-	-	-	-
65	1787	-	-	-	-
66	757	-	-	-	-

Surprisingly, the compounds of this invention selectively modulate the glucocorticoid receptor in relation to the progesterone receptor, mineralocorticoid receptor, androgen receptor, and estrogen receptor-alpha. Therefore these compounds are useful for the treatment of inflammatory, immune, adrenal imbalance, cognitive and behavioral diseases.

The present invention also provides pharmaceutical compositions which comprise compounds of the present invention formulated together with one or more non-toxic pharmaceutically acceptable carriers. The pharmaceutical compositions may be specially formulated for oral administration in solid or liquid form, for parenteral injection, or for rectal administration.

The pharmaceutical compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, or as an oral or nasal spray. The term "parenteral" administration refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Pharmaceutical compositions of this invention for parenteral injection comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required

particle size in the case of dispersions, and by the use of surfactants. Conversely, reduced particle size may maintain biological activity.

These compositions may also contain adjuvants such as preservative, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of the drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption

accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the
5 dosage form may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

The solid dosage forms of tablets, dragees, capsules, pills, and granules can be
10 prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

15 The active compounds can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example,
20 water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

25 Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, and
30 mixtures thereof.

Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room

temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

Compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic.

Methods to form liposomes are known in the art. See, for example, Prescott, Ed., Methods in Cell Biology, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 et seq.

Dosage forms for topical administration of a compound of this invention include powders, sprays, ointments and inhalants. The active compound is mixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives, buffers, or propellants which may be required. Ophthalmic formulations, eye ointments, powders and solutions are also contemplated as being within the scope of this invention.

Actual dosage levels of active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active compound(s) that is effective to achieve the desired therapeutic response for a particular patient, compositions, and mode of administration. The selected dosage level will depend upon the activity of the particular compound, the route of administration, the severity of the condition being treated, and the condition and prior medical history of the patient being treated. However, it is within the skill of the art to start doses of the compound at levels lower than required for to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

Generally dosage levels of about 1 to about 50, more preferably of about 5 to about 20 mg of active compound per kilogram of body weight per day are administered orally to a mammalian patient. If desired, the effective daily dose may be divided into multiple doses for purposes of administration, e.g. two to four separate doses per day.

Abbreviations

Abbreviations that have been used in the descriptions of the scheme and the examples that follow are: $\text{BF}_3\cdot\text{OEt}_2$ for boron trifluoride diethyl etherate; DMF for N,N-dimethylformamide, DMSO for dimethylsulfoxide; and THF for tetrahydrofuran.

5


Synthetic Methods

The compounds and processes of the present invention will be better understood in connection with the following synthetic schemes which illustrate the methods by which the compounds of the invention can be prepared.

10

Syntheses of the compounds of the present invention are described in Scheme 1 and Scheme 2.


Scheme 1



As exemplified in Scheme 1, benzaldehydes 1 can be treated with two equivalents of an aniline or other electron-rich aromatic compound in the presence of Lewis acids such as aluminum chloride to afford symmetric triarylmethanes A. Under similar condition using one equivalent of aniline or electron-rich aromatic compound at much lower reaction temperature, benzhydryl alcohols B could be formed. Benzhydryl alcohols B were then treated with a different aromatic nucleophile such as another aniline in the presence of Lewis acid such as aluminum

trichloride to afford triarylmethanes C. Alcohols B could also be condensed with phenols using the conditions of the Mitsunobu reaction with reagents such as tributylphosphine and diethylazodicarboxylate to form phenyl ethers D. Treatment of 1B with thiophenols in the presence of protic acids such as p-toluenesulfonic acid as catalyst also afforded phenyl thioethers E. Deprotonation of B with base followed by quenching with aromatic isocyanates provided aminophenyl analogs F.

Scheme 2



As exemplified in Scheme 2, benzophenones **G** can be treated with NaBH_4 or other reducing reagents to form benzhydrols **H**, which were then treated with electron-rich aromatics such as anilines in the presence of POCl_3 or another Lewis acid to afford triarylmethanes **I**.
Alternatively, benzophenones **G** were treated with grignard reagents to form carbinols **J**, which were reduced with agents such as formic acid to afford triarylmethanes **K**.

The compounds and processes of the present invention will be better understood in connection with the following examples, which are intended as an illustration of and not a limitation upon the scope of the invention as defined in the appended claims.

Example 1

4',4'' bis(dimethylamino)-2-chloro-5-nitrotriphenylmethane

A solution of 2-chloro-5-nitrobenzaldehyde (17.28 g, 93.1 mmol) and N,N-dimethylaniline (24.78 g, 205 mmol) in CH₂Cl₂ (200 mL) at 0 °C was treated with AlCl₃ (13.60 g, 102 mmol) in one portion. The mixture was allowed to warm to ambient temperature overnight and quenched with aqueous 1M NaOH. The organic phase was washed with brine and dried (Na₂SO₄), filtered, and concentrated. The crude solid was purified by trituration with 5:1 ethyl acetate/ CH₂Cl₂ to provide the title compound.

mp 188-190 °C;

¹H NMR (300 MHz, DMSO-d₆) δ 8.11 (dd, 1H), 7.74 (dd, 2H), 6.87 (d, 4H), 6.68 (d, 4H), 5.70 (s, 1H), 2.87 (s, 12H);

MS (DCI/NH₃) m/e 410 (M+H)⁺;

Anal. calc'd for C₂₃H₂₄ClN₃O₂: C, 65.94; H, 6.01; N, 10.03. Found: C, 65.86; H, 5.80; N, 9.96.

15

Example 24',4'' bis(dimethylamino)-4-chloro-3-nitrotriphenylmethane

Using the procedure described for Example 1, 4-chloro-3-nitrobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

¹H NMR (300 MHz, DMSO-d₆) δ 7.60 (d, J=2.7 Hz, 1H), 7.39 (s, J=7.8 Hz, 1H), 7.27 (dd, J=2.7, 7.8 Hz, 1H), 6.93 (d, J=9.0 Hz, 4H), 6.65 (d, J= 9.0 Hz, 4H), 5.37 (s, 1H), 2.93 (s, 8H);

MS (DCI/NH₃) m/e 410 (M+H)⁺.

25

Example 34',4'' bis(dimethylamino)-5-acetamido-2-chlorotriphenylmethane

Using the procedure described for Example 1, 5-acetamido-2-chlorobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

¹H NMR (300 MHz, DMSO-d₆) δ 7.75 (dd, J=2.7, 9.0 Hz, 1H), 7.28 (m, 3H), 6.95 (m, 5H), 6.63 (d, J=8.7 Hz, 4H), 5.71 (s, 1H), 2.94 (s, 12H), 2.08 (s, 3H);

MS (DCI/NH₃) m/e 422 (M+H)⁺;

Anal. calc'd for C₂₅H₂₈ClN₃O: C, 71.16; H, 6.69; N, 9.96. Found: C, 71.81; H, 6.76; N, 9.96.

Example 4

4',4'' bis(dimethylamino)-4-nitrotriphenylmethane

Using the procedure described for Example 1, 4-nitrobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 181-183 °C;

5 ¹H NMR (300 MHz, DMSO-d₆) δ 8.11 (d, J=9.0 Hz, 2H), 7.27 (d, J=9.0 Hz, 2H), 6.94 (d, J=8.7 Hz, 4H), 6.68 (d, J=8.7 Hz, 4H), 5.43 (s, 1H), 2.44 (s, 12H);

MS (DCI/NH₃) m/e 376 (M+H)⁺;

Anal. calc'd for C₂₃H₂₅N₃O₂: C, 73.58; H, 6.71 ; N, 11.19. Found: C, 73.55; H, 6.75; N, 11.16.

10

Example 5 4',4'' bis(dimethylamino)-4-chlorotriphenylmethane

Using the procedure described for Example 1, 4-chlorobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 98-100 °C;

15 ¹H NMR (300 MHz, DMSO-d₆) δ 2.89 (s, 12H), 5.35 (s, 1H), 6.67 (dd, 4H), 6.94 (dd, 4H), 7.11 (dd, 2H), 7.28 (dd, 2H);

MS (DCI/NH₃) m/e 365 (M+H)⁺;

Anal. calc'd for C₂₃H₂₅ClN₂: C, 75.70; H, 6.90 ; N, 7.67. Found: C, 75.76; H, 6.91; N, 7.54.

20

Example 64',4'' bis(dimethylamino)-3-chlorotriphenylmethane

Using the procedure described for Example 1, 3-chlorobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

25 mp 102-104 °C;

¹H NMR (300 MHz, DMSO-d₆) δ 3.35 (s, 12H), 5.82 (s, 1H), 7.13 (dt, 4H), 7.40 (dt, 4H), 7.51-7.75 (m, 4H);

MS (DCI/NH₃) m/e 365 (M+H)⁺;

Anal. calc'd for C₂₃H₂₅ClN₂: C, 75.70; H, 6.90; N, 7.67. Found: C, 75.76; H, 6.87; N, 7.62.

30

Example 74',4'' bis(dimethylamino)-2-chlorotriphenylmethane

Using the procedure described for Example 1, 2-chlorobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 144-145 °C;

¹H NMR (300 MHz, DMSO-d₆) δ 2.90 (s, 6H), 5.74 (s, 1H), 6.67 (d, J=9 Hz, 4H), 6.89 (d, J=9 Hz, 4H), 7.03 (m, 1H), 7.22 (m, 2H), 7.37 (m, 1H);

MS (DCI/NH₃) m/e 382 (M+ NH₄)⁺, 365 (M+H)⁺;

Anal. calc'd for C₂₃H₂₅ClN₂: C, 75.70; H, 6.90; N, 7.67. Found: C, 75.76; H, 6.81; N, 7.57.

10

Example 8

4',4'' bis(dimethylamino)-2-methoxytriphenylmethane

Using the procedure described for Example 1, 2-methoxybenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 151-153 °C;

15 MS (DCI/NH₃) m/e 361 (M+H)⁺;

Anal. calc'd for C₂₄H₂₈N₂O: C, 79.96; H, 7.82; N, 7.77. Found: C, 79.85; H, 7.71; N, 7.70.

20

Example 9 4',4'' bis(dimethylamino)-3-nitrotriphenylmethane

Using the procedure described for Example 1, 3-nitrobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 141-142 °C;

¹H NMR (300 MHz, DMSO-d₆) δ 8.08 (dt, J=6.3, 2.7 Hz, 1H), 7.90 (m, 1H), 7.58 (m, 2H), 6.91 (d, J=9.0 Hz, 4H), 6.63 (d, J=9.0 Hz, 4H), 5.54 (s, 1H), 2.83 (s, 12H);

25 MS (DCI/NH₃) m/e 376 (M+H)⁺;

Anal. calc'd for C₂₃H₂₅N₃O₂: C, 72.70; H, 6.76; N, 11.05. Found: C, 72.83; H, 6.62; N, 11.02.

30

Example 10

4',4'' bis(dimethylamino)-2-trifluoromethyltriphenylmethane

Using the procedure described for Example 1, 2-trifluoromethylbenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 94-96 °C;

Anal. calc'd for $C_{24}H_{25}F_3N_2$: C, 72.34; H, 6.32; N, 7.03. Found: C, 72.12; H, 6.29; N, 7.00.

Example 11

5

4',4'' bis(dimethylamino)triphenylmethane

Using the procedure described for Example 1, benzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.
mp 100-102 °C.

10

Example 12

4',4'' bis(dimethylamino)-2-chloro-6-nitrotriphenylmethane

Using the procedure described for Example 1, 2-chloro-6-nitrobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

1H NMR (300 MHz, DMSO- d_6) δ 7.57 (dd, J=1.5, 8.4 Hz, 1H), 7.43 (dd, J=1.5, 8.4 Hz, 1H), 7.28 (t, J=8.4 Hz, 1H), 6.98 (d, J=9.0 Hz, 4H), 6.64 (d, J=9.0 Hz, 4H), 6.02 (s, 1H), 2.42 (s, 12H);

15

MS (DCI/ NH_3) m/e 410 (M+H) $^+$;

Anal. calc'd for $C_{23}H_{24}ClO_2N_3$: C, 67.39; H, 5.9; N, 10.25. Found: C, 67.29; H, 5.87; N, 10.07.

20

Example 13

4',4'' bis(N-piperidinyl)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 1, 2-chloro-5-nitrobenzaldehyde and N-phenylpiperidine were treated with a Lewis acid to provide the title compound.

1H NMR (300 MHz, DMSO- d_6) δ 8.02 (dd, J=3.0, 9.0 Hz, 1H), 7.88 (d, J=3.0 Hz, 1H), 7.50 (d, J=9.0 Hz, 1H), 6.91 (d, J=9.0 Hz, 4H), 6.83 (d, J=9.0 Hz, 4H), 5.77 (s, 1H), 3.15 (t, J = 5.4 Hz, 8H), 1.70 (m, 8H), 1.55 (m, 4H);

25

MS (DCI/ NH_3) m/e 490 (M+H) $^+$;

Anal. calc'd for $C_{29}H_{32}N_3ClO_2$: C, 71.08; H, 6.56; N, 8.57. Found: C, 70.94; H, 6.48; N, 8.47.

30

Example 14

4',4'' bis(dimethylamino)-2-bromotriphenylmethane

Using the procedure described for Example 1, 2-bromobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 152-153 °C;

Anal. calc'd for $C_{23}H_{25}BrN_2$: C, 67.48; H, 6.15; N, 6.84. Found: C, 67.28; H, 6.21; N, 6.84.

Example 15

4',4'' bis(dimethylamino)-2-methyltriphenylmethane

Using the procedure described for Example 1, o-tolualdehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 98-100 °C;

Anal. calc'd for $C_{24}H_{28}N_2$: C, 83.67; H, 8.19; N, 8.13. Found: C, 83.72; H, 8.31; N, 8.07.

Example 16

4',4'' bis(dimethylamino)-2,3,5-trichlorotriphenylmethane

Using the procedure described for Example 1, 2,3,5-trichlorobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 149-151 °C;

Anal. calc'd for $C_{23}H_{23}Cl_3N_2$: C, 63.68; H, 5.34; N, 6.45. Found: C, 63.39; H, 5.11; N, 6.35.

Example 17

4',4'' bis(dimethylamino)-2-chloro-5-trifluoromethyltriphenylmethane

Using the procedure described for Example 1, 2-chloro-5-trifluoromethylbenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 143-145 °C;

Anal. calc'd for $C_{24}H_{24}ClF_3N_2$: C, 66.58; H, 5.58; N, 6.47. Found: C, 66.40; H, 5.78; N, 6.37.

Example 18

4',4'' bis(dimethylamino)-2,4-dichlorotriphenylmethane

Using the procedure described for Example 1, 2,4-dichlorobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 104-105 °C;

Anal. calc'd for $C_{23}H_{24}Cl_2N_2$: C, 69.17; H, 6.05; N, 7.01. Found: C, 69.17; H, 5.83; N, 6.90.

Example 19

5 4',4" bis(dimethylamino)-2-chloro-4,5-methylenedioxytriphenylmethane

Using the procedure described for Example 1, 2-chloro-4,5-methylenedioxybenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 168-170 °C;

10 Anal. calc'd for $C_{24}H_{25}ClN_2O_2$: C, 70.49; H, 6.16; N, 6.85. Found: C, 70.19; H, 5.99 ; N, 6.63.

Example 20

15 4',4" bis(dimethylamino)-2,6-dichlorotriphenylmethane

Using the procedure described for Example 1, 2,6-dichlorobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 134-136 °C;

Anal. calc'd for $C_{23}H_{24}Cl_2N_2$: C, 69.17; H, 6.05; N, 7.01. Found: C, 68.95; H, 5.93; N, 6.81.

20

Example 21

25 4',4" bis(dimethylamino)-2,3-dichlorotriphenylmethane

Using the procedure described for Example 1, 2,3-dichlorobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 172-174 °C;

Anal. calc'd for $C_{23}H_{24}Cl_2N_2$: C, 69.17; H, 6.05 ; N, 7.01. Found: C, 69.08; H, 5.98; N, 6.89.

Example 22

30 4',4" bis(N-morpholinyl)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 1, 2-chloro-5-nitrobenzaldehyde and N-phenyl morpholine were treated with a Lewis acid to provide the title compound.

mp 126-128 °C;

¹H NMR (300 MHz, DMSO-d₆) δ 8.02 (dd, J=3.0, 9.0 Hz, 1H), 7.88 (d, J=3.0 Hz, 1H), 7.53 (d, J=9.0 Hz, 1H), 6.95 (d, J=8.8 Hz, 4H), 6.83 (d, J=8.8 Hz, 4H), 3.88 (t, J=4.5 Hz, 8H), 3.15 (t, J=4.5 Hz, 8H);

MS (DCI/NH₃) m/e 494 (M+H)⁺;

- 5 Anal. calc'd for C₂₇H₂₈N₃O₄Cl: C, 65.65; H, 5.71; N, 8.51. Found: C, 65.49; H, 5.65; N, 8.24.

Example 23

4',4'' bis(methylamino)-2-chloro-5-nitrotriphenylmethane

- 10 Using the procedure described for Example 1, 2-chloro-5-nitrobenzaldehyde and N-methyl aniline were treated with a Lewis acid to provide the title compound.

¹H NMR (300 MHz, DMSO-d₆) δ 8.01 (dd, J=3.0, 9.0 Hz, 1H), 8.89 (d, J=3.0 Hz, 1H), 7.53 (d, J=9.0 Hz, 1H), 6.88 (d, J=8.8 Hz, 4H), 6.55 (d, J=8.8 Hz, 4H), 5.75 (s, 1H), 3.70 (brs, 2H), 2.84 (s, 12H);

- 15 MS (DCI/NH₃) m/e 382 (M+H)⁺, 399 (M+NH₄)⁺;

Anal. calc'd for C₂₁H₂₀N₃O₂Cl: C, 66.05; H, 5.28; N, 11.00. Found: C, 66.01; H, 5.09; N, 10.63.

Example 24

4',4'' bis(dimethylamino)-2,5-dichlorotriphenylmethane

- 20 Using the procedure described for Example 1, 2,5-dichlorobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 181-183 °C;

Anal. calc'd for C₂₃H₂₄Cl₂N₂: C, 69.17; H, 6.05; N, 7.01. Found: C, 69.23; H, 6.05; N,

- 25 7.00.

Example 25

4',4'' bis(dimethylamino)-2-fluorotriphenylmethane

- 30 Using the procedure described for Example 1, 2-fluorobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 129-131 °C;

Anal. calc'd for C₂₃H₂₅FN₂: C, 79.27; H, 7.23; N, 8.03. Found: C, 79.32; H, 7.17; N, 7.96.

Example 26

4',4'' bis(dimethylamino)-2-iodotriphenylmethane

Using the procedure described for Example 1, 2-iodobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 143-145 °C;

- 5 Anal. calc'd for $C_{23}H_{25}IN_2$: C, 60.53; H, 5.52; N, 6.13 . Found: C, 60.68; H, 5.54; N, 6.17.

Example 274',4'' bis(methylamino)-2,5-dichlorotriphenylmethane

- 10 Using the procedure described for Example 1, 2,5-dichlorobenzaldehyde and N-methyl aniline were treated with a Lewis acid to provide the title compound.

mp 135-140 °C;

1H NMR (300 MHz, DMSO- d_6) δ 7.48 (d, J=9.0 Hz, 1H), 7.31 (dd, J=3.0, 9.0 Hz, 1H), 6.87 (d, J=3.0 Hz, 1H), 6.76 (d, J=8.8 Hz, 4H), 6.45 (d, J=8.8 Hz, 4H), 5.58 (brs, 2H), 5.55 (s, 1H), 2.65 (s, 6H);

- 15 MS (DCI/ NH_3) m/e 371 (M+H) $^+$;

Anal. calc'd for $C_{21}H_{20}N_2Cl_2$: C, 68.09; H, 5.45 ; N, 7.57. Found: C, 67.83; H, 5.19; N, 7.52.

Example 28

- 20 4',4'' bis(N-morpholinyl)-2,3,5-trichlorotriphenylmethane

Using the procedure described for Example 1, 2,3,5-trichlorobenzaldehyde and N-phenyl morpholine were treated with a Lewis acid to provide the title compound.

mp 102-104 °C;

- 25 Anal. calc'd for $C_{27}H_{27}N_2O_2Cl_3$: C, 62.78; H, 5.27; N, 5.43. Found: C, 63.19; H, 5.16; N, 4.99.

Example 294',4'' bis(N-pyrrolidinyl)-2-chloro-5-nitrotriphenylmethane

- 30 Using the procedure described for Example 1, 2-chloro-5-nitrobenzaldehyde and N-phenyl pyrrolidine were treated with a Lewis acid to provide the title compound.

mp > 200 °C;

Anal. calc'd for $C_{27}H_{28}ClN_3O_2$: C, 70.19; H, 6.10; N, 9.09. Found: C, 70.45; H, 6.03; N, 8.83.

Example 304',4'' bis(di-n-butylamino)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 1, 2-chloro-5-nitrobenzaldehyde and N,N-dibutyl aniline were treated with a Lewis acid to provide the title compound.

5 mp 69-71 °C;

Anal. calc'd for $C_{35}H_{48}ClN_3O_2$: C, 72.70; H, 8.36; N, 7.26. Found: C, 73.04; H, 8.62; N, 7.32.

Example 31

10 4',4'' bis(N-(2-acetoxyethyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 1, 2-chloro-5-nitrobenzaldehyde and N-(2-acetoxyethyl)-N-methyl aniline were treated with a Lewis acid to provide the title compound.

mp 100-102 °C;

Anal. calc'd for $C_{29}H_{32}ClN_3O_6$: C, 62.86; H, 5.82; N, 7.58. Found: C, 62.92; H, 5.79; N, 7.49.

Example 324',4'' bis(N-(2-hydroxyethyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 1, 2-chloro-5-nitrobenzaldehyde and N-(2-hydroxyethyl)-N-methyl aniline were treated with a Lewis acid to provide the title compound.

20 mp 150-152 °C;

Anal. calc'd for $C_{25}H_{28}ClN_3O_4$: C, 63.89; H, 6.00; N, 8.94. Found: C, 63.72; H, 5.76; N, 8.72.

Example 334',4'' bis(dimethylamino)-2-chloro-5-iodotriphenylmethane

Using the procedure described for Example 1, 2-chloro-5-iodobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

Anal. calc'd for $C_{23}H_{24}N_2ClI$: C, 56.28; H, 4.93; N, 5.71. Found: C, 56.23; H, 4.79; N, 5.58.

Example 344',4'' bis(dimethylamino)-5-bromo-2-chlorotriphenylmethane

Using the procedure described for Example 1, 5-bromo-2-chlorobenzaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 177-178 °C;

¹H NMR (300 MHz, DMSO-d₆) δ 7.26 (dd, J=2.7, 8.3 Hz, 1H), 7.20 (d, J=8.3 Hz, 1H),

5 7.12 (d, J=2.7 Hz, 1H), 6.92 (d, J=9.0 Hz, 4H), 6.67 (d, J=9.0 Hz, 4H), 5.70 (s, 1H), 3.92 (s, 12 H);

MS (DCI/NH₃) m/e 445 (M+H)⁺;

Anal. calc'd for C₂₃H₂₄N₂ClBr: C, 62.25; H, 5.45; N, 6.31. Found: C, 62.06; H, 5.24; N, 6.18.

10

Example 35

4',4" bis(N-(t-butoxycarbonyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 1, 2-chloro-5-nitrobenzaldehyde and N-(t-butoxycarbonyl) N-methyl aniline were treated with a Lewis acid to provide the title compound.

15 mp 180-183 °C;

¹H NMR (300 MHz, DMSO-d₆) δ 8.08 (dd, J=3.0, 8.7 Hz, 1H), 7.86 (d, J=3.0 Hz, 1H),

7.55 (d, J=8.7 Hz, 1H), 7.20 (d, J=8.7 Hz, 4H), 7.04 (d, J=8.7 Hz, 4H), 5.92 (s, 1H), 3.27 (s, 6H), 1.47 (s, 19H);

MS (DCI/NH₃) m/e 599 (M+H)⁺.

20

Example 36

4',4" bis(N-benzylamino)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 1, 2-chloro-5-nitrobenzaldehyde and N-benzyl aniline were treated with a Lewis acid to provide the title compound.

25 mp 158-160 °C;

¹H NMR (300 MHz, DMSO-d₆) δ 8.00 (dd, J=3.0, 8.8 Hz, 1H), 7.92 (d, J=3.0 Hz, 1H),

7.52 (d, J=8.8 Hz, 1H), 7.3 (m, 10H), 6.85 (d, J=9.0 Hz, 4H), 6.59 (d, J=9.0 Hz, 4H), 5.74 (s, 1H), 4.30 (s, 4H), 4.08 (brs, 2H);

MS (DCI/NH₃) m/e 551 (M+NH₄)⁺, 534 (M+H)⁺;

30 Anal. calc'd for C₃₃H₂₆N₃O₂Cl: C, 74.50; H, 4.93; N, 7.01. Found: C, 74.60; H, 4.95; N, 6.70.

Example 37

4',4" bis(N-benzylamino)-2,5-dichlorotriphenylmethane

Using the procedure described for Example 1, 2,5-dichlorobenzaldehyde and N-benzyl aniline were treated with a Lewis acid to provide the title compound.

¹H NMR (300 MHz, DMSO-d₆) δ 7.30 (m, 1H), 7.11 (dd, J=3.0, 8.8 Hz, 1H), 6.97 (d, J=3.0 Hz, 1H), 6.85 (d, J=9.0 Hz, 4H), 6.59 (d, J=9.0 Hz, 4H), 6.59 (s, 1H), 4.30 (s, 4H), 4.02 (BRS. 2H);

MS (DCI/NH₃) m/e 523 (M+H)⁺;

Anal. calc'd for C₃₃H₂₈N₂Cl₂: C, 75.71; H, 5.39; N, 5.35. Found: C, 75.41; H, 5.24; N, 5.17.

Example 38

4',4'' bis(dimethylamino)-4-methoxytriphenylmethane

Using the procedure described for Example 1, p-anisaldehyde and N,N-dimethylaniline were treated with a Lewis acid to provide the title compound.

mp 101-103 °C;

Anal. calc'd for C₂₄H₂₈N₂O: C, 79.96; H, 7.82; N, 7.77. Found: C, 79.90; H, 7.76; N, 7.74.

Example 39

4'-dimethylamino-4''-methylamino-2-chloro-5-nitrotriphenylmethane

Example 39A

4'-dimethylamino-2-chloro-5-nitrodiphenylmethanol

A solution of 2-chloro-5-nitrobenzaldehyde (2.09 g, 11 mmol) and N,N-dimethylaniline (1.21 g, 10 mmol) in CH₂Cl₂ (50 ml) at -78 °C was treated with solid AlCl₃ (1.33 g, 10 mmol) portionwise. The mixture was kept at this temperature for 4 hours and quenched with aqueous 1M NaOH. The organic phase was washed with brine, dried (Na₂SO₄), filtered, and concentrated to dryness. The residue was flash chromatographed on silica gel with 10-20% ethyl acetate/hexane to provide the title compound
Method C.

Example 39B

4'-dimethylamino-4''-methylamino-2-chloro-5-nitrotriphenylmethane

A solution of Example 39A (0.31 g, 1 mmol) and N-methylaniline (0.21 g, 2 mmol) in CH₂Cl₂ (15 ml) at 0 °C was treated with solid AlCl₃ (0.40 g, 3 mmol) portionwise. The

mixture was allowed to warm to ambient temperature overnight and quenched with aqueous 1M NaOH. The organic phase was washed with brine, dried (Na_2SO_4), filtered, and concentrated. The residue was flash chromatographed on silica gel with 10-20% ethyl acetate/hexane to provide the title compound.

5 mp 152-153 °C;

Anal. calc'd for $\text{C}_{22}\text{H}_{22}\text{ClN}_3\text{O}_2$: C, 66.74; H, 5.60; N, 10.61. Found: C, 66.74; H, 5.70; N, 10.68.

Example 40

10 4'-dimethylamino-4''-(N-morpholinyl)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 39B, a solution of Example 39A was treated with N-phenyl morpholine and a Lewis acid to provide the title compound.

mp 194-196 °C;

Anal. calc'd for $\text{C}_{25}\text{H}_{26}\text{ClN}_3\text{O}_3$: C, 66.43; H, 5.79; N, 9.29. Found: C, 66.52; H, 5.43; N, 15 9.19.

Example 41

4'-dimethylamino-4''-(N-pyrrolidinyl)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 39B, a solution of Example 39A was treated 20 with N-phenyl pyrrolidine and a Lewis acid to provide the title compound.

mp 157-158 °C;

Anal. calc'd for $\text{C}_{25}\text{H}_{26}\text{ClN}_3\text{O}_2$: C, 68.87; H, 6.01; N, 9.63. Found: C, 68.92; H, 6.02; N, 9.65.

25

Example 42

4'-dimethylamino-4''-(N-methyl-N-(2-hydroxyethyl)amino)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 39B, a solution of Example 39A was treated with N-(2-hydroxyethyl)-N-methyl aniline and a Lewis acid to provide the title compound.

mp 136-148 °C;

30 Anal. calc'd for $\text{C}_{24}\text{H}_{26}\text{ClN}_3\text{O}_3$: C, 65.52; H, 5.95; N, 9.55. Found: C, 65.50; H, 5.86; N, 9.64.

Example 43

4'-dimethylamino-4''-(di-n-butylamino)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 39B, a solution of Example 39A was treated with dibutylaniline and a Lewis acid to provide the title compound.

mp 115-118 °C;

Anal. calc'd for $C_{29}H_{36}ClN_3O_2$: C, 70.49; H, 7.34; N, 8.50. Found: C, 70.40; H, 7.34; N, 8.58.

Example 44

3'-methyl-4',4''-bis(dimethylamino)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 39B, a solution of Example 39A was treated with N,N,2-trimethylaniline and a Lewis acid to provide the title compound.

mp 174-176 °C;

Anal. calc'd for $C_{24}H_{26}ClN_3O_2$: C, 67.89; H, 6.18; N, 9.91. Found: C, 67.73; H, 5.82; N, 10.08.

Example 46

4'-dimethylamino-4''-(N-piperidinyl)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 39B, a solution of Example 39A was treated with N-phenyl piperidine and a Lewis acid to provide the title compound.

mp 166-168 °C;

Anal. calc'd for $C_{26}H_{28}ClN_3O_2$: C, 69.40; H, 6.27; N, 9.33. Found: C, 69.44; H, 6.19; N, 9.20.

Example 48

4'-methylamino-4''-(t-butoxycarbonylamino)-2-chloro-5-nitrotriphenylmethane

Example 48A

4'-methylamino-2-chloro-5-nitrodiphenylmethanol

Using the procedure described for Example 39A, 2-chloro-5-nitrobenzaldehyde was treated with N-methyl aniline to provide the title compound.

Example 48B

Using the procedure described for Example 39B, a solution of Example 48A was treated with N-methyl-N-(t-butoxycarbonyl) aniline and a Lewis acid to provide the title compound.

mp 141-144 °C;

¹H NMR (300 MHz, DMSO-d₆) δ 8.04 (dd, J=3.0, 9.0 Hz, 1H), 7.88 (d, J=3.0 Hz, 1H), 7.55 (d, J=9.0 Hz, 1H), 7.18 (d, J=8.7 Hz, 2H), 7.02 (d, J=8.7 Hz, 2H), 6.88 (d, J=9.0 Hz, 2H), 6.54 (d, J=9.0 Hz, 2H), 5.42 (s, 1H), 3.72 (brs, 1H), 3.27 (s, 3H), 2.32 (s, 3H), 1.48 (s, 9H);

5 MS (DCI/NH₃) m/e 499 (M+NH₄)⁺, 481 (M+H)⁺.

Example 49

4'-methylamino-4''-(N-(2-acetoxyethyl)-N-methylamino)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 39B, a solution of Example 48A was treated
10 with N-methyl-N-(2-acetoxyethyl) aniline and a Lewis acid to provide the title compound.
mp 117-119 °C;

Anal. calc'd for C₂₅H₂₆ClN₃O₄: C, 64.16; H, 5.60; N, 8.97. Found: C, 64.22; H, 5.64; N, 8.91.

15

Example 50

4'-dimethylamino-4''-(N-methyl-N-(2-benzoyloxyethyl)amino)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 60, a solution of Example 42 was treated
with benzoyl chloride in dichloromethane to provide the title compound.

20 mp 143-145 °C;

Anal. calc'd for C₃₁H₃₀ClN₃O₄: C, 68.43; H, 5.55; N, 7.72. Found: C, 68.79; H, 5.62; N, 7.60.

Example 51

4'-dimethylamino-4''-(N-methyl-N-(2-acetoxyethyl)amino)-2-chloro-5-nitrotriphenylmethane

Using the procedure described for Example 60, a solution of Example 42 was treated
with acetyl chloride in dichloromethane to provide the title compound.

Example 52

phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] ether

30 A solution of Example 39A (0.31 g, 1 mmol), phenol (0.10 g, 1.1 mmol) and triphenylphosphine (0.39 g, 1.5 mmol) in THF (10 ml) at 0 °C was treated with diethylazodicarboxylate (0.27 g, 1.5 mmol). The final solution was allowed to warm to ambient temperature overnight and quenched with H₂O and extracted with ethyl acetate. The

organic phase was washed with brine and dried (Na_2SO_4), filtered, and concentrated to dryness. The crude viscous oil was purified by flash chromatography on silica gel with 20-30% ethyl acetate/hexane to provide the title compound.

mp 126-128 °C;

- 5 Anal. calc'd for $\text{C}_{21}\text{H}_{19}\text{ClN}_2\text{O}_3$: C, 65.88; H, 5.00; N, 7.31. Found: C, 65.50; H, 4.89; N, 7.11.

Example 53

4-chlorophenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] ether

- 10 Using the procedure described for Example 52, a solution of Example 39A was treated with 4-chloro phenol to provide the title compound.

mp 135-137 °C;

Anal. calc'd for $\text{C}_{21}\text{H}_{18}\text{Cl}_2\text{N}_2\text{O}_3$: C, 60.44; H, 4.34; N, 6.71. Found: C, 60.10; H, 4.24; N, 6.56.

15

Example 54

phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] thioether

A solution of Example 39A (0.31 g, 1 mmol), thiophenol (1.10 g, 10 mmol) and p-toluenesulfonic acid (19 mg, 0.1 mmol) in toluene (10 ml) was heated to reflux for 4 hours.

- 20 After cooling, the solution was washed with 1M NaOH (aq) and brine respectively, dried (Na_2SO_4), filtered, and concentrated. The crude solid was purified by flash chromatography on silica gel with 10-20% ethyl acetate/hexane to provide the title compound.

mp 134-136 °C;

- 25 Anal. calc'd for $\text{C}_{21}\text{H}_{19}\text{ClN}_2\text{O}_2\text{S}$: C, 63.23; H, 4.80; N, 7.02. Found: C, 63.08; H, 4.54; N, 6.85.

Example 55

phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] amine

- 30 A solution of Example 39A (0.31 g, 1 mmol) and phenylisocyanate (0.14 g, 1.2 mmol) in THF (10 ml) at 0 °C was treated with potassium t-butoxide (1M in THF, 0.11 ml, 1.1 mmol). The resulting solution was allowed to warm to ambient temperature overnight, quenched with H_2O , and extracted with ethyl acetate. The organic phase was washed with brine and dried (Na_2SO_4), filtered, and concentrated. The crude solid was purified by recrystallization from ethanol to provide the title compound.

mp 195-197 °C;

Anal. calc'd for $C_{21}H_{20}ClN_3O_2$: C, 65.28; H, 5.34; N, 10.87. Found: C, 65.53; H, 5.11; N, 10.79.

5

Example 56

4'-dimethylamino-2-chloro-5-nitrotriphenylmethane

Example 56A

2-chloro-5-nitrodiphenylmethanol

- 10 To a solution of 2-chloro-5-nitrobenzophenone (2.62 g, 10 mmol) in methanol (20 ml) at 0°C was added solid $NaBH_4$ (170 mg, 4.5 mmol) in portions. The mixture was allowed to warm to ambient temperature for 4 hours, quenched with saturated NH_4Cl (aq) and extracted with ethyl acetate. The organic phase was washed with brine and dried (Na_2SO_4), filtered, and concentrated. The crude solid was purified by flash column chromatography with 1:5 ethyl
15 acetate/hexanes to provide the title compound.
MS (DCI/ NH_3) m/e 264 (M+H)⁺.

Example 56B

4'-dimethylamino-2-chloro-5-nitrotriphenylmethane

- 20 A solution of Example 56A (0.66 g, 2.50 mmol) and $POCl_3$ in N,N-dimethylaniline (15 ml) was heated to 100 °C for 12 hours. After cooling to ambient temperature, the solution was washed with 1M $NaOH$ (aq) and brine respectively and dried (Na_2SO_4), filtered, and concentrated. The crude solid was purified by flash column chromatography with 1:5 ethyl acetate/hexanes to provide the title compound.
- 25 mp 127-129 °C;
 1H NMR (300 MHz, $DMSO-d_6$) δ 8.13 (dd, 1H), 7.87 (d, 1H), 7.74 (d, 1H), 7.37-7.23 (m, 3H), 7.13 (d, 2H), 6.96 (d, 2H), 6.73 (d, 2H), 5.92 (s, 1H), 2.93 (s, 6H);
MS (DCI/ NH_3) m/e 367 (M+H)⁺;
Anal. calc'd for $C_{21}H_{19}ClN_2O_2$: 68.75; H, 5.22; N, 7.63. Found: C, 68.87; H, 5.18; N,
30 7.60.

Example 57

4'-dimethylamino-2-chlorotriphenylmethanol

A solution of N,N-dimethyl-4-bromoaniline (2.00 g, 10 mmol) and Mg (0.24 g, 10 mmol) in THF (40 ml) was heated to reflux until it became a clear solution. To this ice-cooled solution was added 2-chlorobenzophenone (2.17 g, 10 mmol) in portions. The resulting solution was allowed to warm to ambient temperature for 12 hours, quenched with saturated NH₄Cl (aq) and extracted with ethyl acetate. The organic phase was washed with brine and dried (Na₂SO₄), filtered, and concentrated. The crude solid was purified by flash column chromatography with 1:5 ethyl acetate/hexanes to provide the title compound.
mp 130-133 °C;
¹H NMR (300 Mhz, acetone-d₆) δ 7.40-7.19 (m, 8H), 7.08-7.02 (m, 3H), 6.66 (d, 2H), 2.94 (s, 6H), 2.78 (s, 1H);
MS (DCI/NH₃) m/e 338 (M+H)⁺;
Anal. calc'd for C₂₁H₂₀ClNO: C, 74.65; H, 5.96; N, 4.14. Found: C, 74.69; H, 5.91; N, 4.15.

15

Example 584'-dimethylamino-2-chlorotriphenylmethane

A solution of Example 57 (0.50 g, 1.48 mmol), saturated Na₂CO₃ (2 ml) and formic acid (10 ml) was heated at reflux for 12 hours. After cooling to ambient temperature, the solution was quenched with saturated Na₂CO₃ and extracted with ethyl acetate. The organic layer was washed with 1M NaOH (aq) and brine respectively and dried (Na₂SO₄), filtered, and concentrated. The residue was further treated with LiAlH₄ (1 M in THF, 5 ml) and quenched with saturated NH₄Cl (aq) followed by another extraction with ethyl acetate. The organic phase was again washed with brine and dried (Na₂SO₄), filtered, and concentrated. The crude solid was purified by flash column chromatography with 1:5 ethyl acetate/hexanes to provide the title compound.
mp 122-124 °C;
¹H NMR (300 Mhz, acetone-d₆) δ 7.40-6.90 (m, 11H), 6.69(dd, 2H), 5.84 (s, 1H), 2.90(6H);
MS (DCI/NH₃) m/e 322 (M+H)⁺.

25
30Example 594',4'' bis(dimethylamino)-5-amino-2-chlorotriphenylmethane

A solution of Example 1 (5.0 g, 12.2 mmol) and Pd/C (5%, 0.5g) was suspended in 200 mL of methanol and degassed under vacuum. A balloon of hydrogen gas was affixed to

the flask and the solution was stirred vigorously for 2 hours. The balloon was removed and the solution was again degassed and then filtered through celite. The solution was concentrated in vacuo, and the crude product was crystallized from hexane:methylene chloride (2:1) to provide the title compound.

- 5 mp 194-197 °C;
1H NMR (300 MHz, DMSO-d₆) δ 6.97 (d, J=8.7 Hz, 1H), 6.88 (d, J=9 Hz, 4H), 6.65 (d, J=9.0 Hz, 4H), 6.49 (dd, J=3.0, 8.7 Hz, 1H), 6.22 (d, J=3.0 Hz, 1H), 5.50 (s, 1H), 5.10 (brs, 2H), 2.45 (s, 12H);
MS (DCI/NH₃) m/e 380 (M+H)⁺;
10 Anal. calc'd for C₂₃H₂₆N₃Cl : C, 72.71; H, 6.90 ; N, 11.06. Found: C, 72.16; H, 6.87; N, 10.66.

Example 60

4',4'' bis(dimethylamino)-2-chloro-5-(4-nitrobenzamido)triphenylmethane

- 15 A solution of Example 59 (0.20 g, 0.53 mmol) in methylene chloride (5 mL) was cooled to 0 °C. Diisopropylethylamine (85 mg, 0.66 mmol) was added followed by 4-nitrobenzoyl chloride (0.11 g, 0.58 mmol). The solution was stirred for 24 hours and filtered through a silica column topped with MgSO₄ using methylene chloride as eluent. The filtrate was concentrated in vacuo, and the crude product was crystallized from chloroform:hexane 1:3 to provide the title compound.
20 1H NMR (300 MHz, DMSO-d₆) δ 8.32 (d, J=8.7 Hz, 2H), 7.98 (d, J=8.7 Hz, 2H), 7.90 (d, J=8.7 Hz, 1H), 7.69 (s, 1H), 7.38 (d, J=8.7 Hz, 1H), 6.95 (d, J=9.0 Hz, 4H), 6.82 (d, J=3.0 Hz, 6.65 (d, J=9.0 Hz, 4H), 5.78 (s, 1H), 2.95 (s, 12H), ppm;
Anal. calc'd for C₃₀H₂₉N₄O₃Cl: C, 68.11; H, 5.53; N, 10.59. Found: C, 68.31; H, 5.45; N, 10.47.
25

Example 61

4',4'' bis(dimethylamino)-2-chloro-5-(4-nitrocinnamido)triphenylmethane

- Using the procedure described for Example 60, a solution of Example 59 was treated
30 with 4-nitro cinnamoyl chloride to provide the title compound.
1H NMR (300 MHz, DMSO-d₆) δ 10.39 (s, 1H), 8.29 (d, J=9.0 Hz, 2H), 7.85 (d, J=9.0 Hz, 2H), 7.81 (dd, J=3.0, 8.7 Hz, 1H), 7.68 (d, J=15.3 Hz, 1H), 7.38 (d, J=8.7 Hz, 1H), 7.23 (d, J=3.0 Hz, 1H), 6.95 (d, J=15.3 Hz, 1H), 6.88 (d, J=9.0 Hz, 4H), 6.67 (d, J=9.0 Hz, 4H), 5.62 (s, 1H), 2.85 (s, 12H);

MS (DCI/NH₃) m/e 555 (M+H)⁺;

Anal. calc'd for C₃₂H₃₁N₄O₃Cl: C, 69.24; H, 5.63; N, 10.09. Found: C, 68.95; H, 5.45; N, 9.80.

5

Example 62

4',4" bis(dimethylamino)-2-chloro-5-(cyclopropylcarbamido)-triphenylmethane

Using the procedure described for Example 60, a solution of Example 59 was treated with cyclopropanecarbonyl chloride to provide the title compound.

¹H NMR (300 MHz, DMSO-d₆) δ 7.82 (d, J=8.7 Hz, 1H), 7.29 (d, J=8.7 Hz, 1H), 7.23 (s, 1H), 6.93 (d, J=9.0 Hz, 4H), 6.68 (d, J=9.0 Hz, 4H), 6.62 (s, 1H), 5.73 (s, 1H), 2.92 (s, 12H), 1.38 (m, 1H), 1.05 (m, 2H), 0.80 (m, 2H);

MS (DCI/NH₃) m/e 448 (M+H)⁺;

Anal. calc'd for C₂₇H₃₀N₃ClO: C, 72.39; H, 6.75; N, 9.38. Found: C, 72.18; H, 6.66; N, 9.25.

15

Example 63

4',4" bis(dimethylamino)-2-chloro-5-(dimethylsulphonimido)triphenylmethane

Using the procedure described for Example 60, a solution of Example 59 was treated with methanesulfonyl chloride to provide the title compound.

¹H NMR (300 MHz, DMSO-d₆) δ 7.43 (d, J=8.8 Hz, 1H), 7.14 (dd, J=3.0, 8.8 Hz, 1H), 6.96 (d, J=3.0 Hz, 1H), 6.93 (d, J=9.0 Hz, 4H), 6.68 (d, J=9.0 Hz, 4H), 3.25 (s, 6H), 2.93 (s, 12H);

MS (DCI/NH₃) m/e 536 (M+H)⁺;

Anal. calc'd for C₂₅H₃₀N₃S₂O₄Cl: C, 56.01; H, 5.64; N, 7.84. Found: C, 55.92; H, 5.64; N, 7.73.

25

Example 64

4',4" bis(dimethylamino)-2-chloro-5-(methoxycarbonylamino)triphenylmethane

Using the procedure described for Example 60, a solution of Example 59 was treated with methyl chloroformate to provide the title compound.

¹H NMR (300 MHz, DMSO-d₆) δ 7.5 (brs, 1H), 7.30 (d, J=8.8 Hz, 1H), 6.94 (d, J=9.0 Hz, 4H), 6.68 (d, J=9.0 Hz, 4H), 6.61 (dd, J=3.0, 8.8 Hz, 1H), 6.45 (s, 1H), 5.73 (s, 1H), 3.73 (s, 3H), 2.94 (s, 12H);

MS (DCI/NH₃) m/e 438 (M+H)⁺;

Anal. calc'd for $C_{25}H_{28}N_3O_2Cl$: C, 68.62; H, 6.45; N, 9.61. Found: C, 68.96; H, 6.70; N, 9.42.

Example 65

5 4',4'' bis(dimethylamino)-2-chloro-5-(2-furanyl)methylimino)triphenylmethane

A solution of Example 59, (0.20g, 0.53 mmol), furfural (63 mg, 0.66 mmol) and p-TsOH·H₂O were dissolved in toluene (2 mL) and heated to 120 °C. The solution was stirred at this temperature for 2 hours followed by cooling to ambient temperature. The solvent was removed in vacuo, and the crude product was crystallized from hexane:methylene chloride (2:1) to provide 0.11 g of the title compound.

¹H NMR (300 MHz, DMSO-d₆) δ 8.38 (s, 1H), 7.89 (d, J=1.0 Hz, 1H), 7.43 (d, J=8.7 Hz, 1H), 7.16 (m, 2H), 6.85 (d, J=9.0 Hz, 4H), 6.77 (d, J=3.0 Hz, 1H), 6.70 (m, 1H), 6.64 (d, J=9.0 Hz, 4H), 5.64 (s, 1H), 2.88 (s, 12H);

MS (DCI/NH₃) m/e 458 (M+H)⁺;

15 Anal. calc'd for $C_{28}H_{28}N_3OCl$: C, 73.43; H, 6.16; N, 9.17. Found: C, 73.31; H, 6.12; N, 8.99.

Example 66

20 4',4'' bis(dimethylamino)-2-chloro-5-(2-furanyl)methylamino)triphenylmethane


A solution of Example 65 (52 mg, 0.11 mmol) and sodium cyanoborohydride (7.1 mg, 0.11 mmol) were suspended in methanol (2 mL) and acetic acid (1 mL) and stirred for 12 hours. Water was added followed by extraction with methylene chloride (10 mL, 3X). The solution was dried (Na₂SO₄), filtered and reduced in vacuo. The crude product was crystallized from hexane:methylene chloride (1:1) to provide 29 mg of the title compound.

25 ¹H NMR (300 MHz, DMSO-d₆) δ 7.33 (d, J=2.1 Hz, 1H), 7.14 (d, J=8.7 Hz, 1H), 6.93 (d, J=9.0 Hz, 4H), 6.68 (d, J=9.0 Hz, 4H), 6.45 (dd, J=3.0, 8.7 Hz, 1H), 6.39 (d, J=6.3 Hz, 6.36 (dd, J=2.1, 6.3 Hz, 1H), 6.08 (d, J= 3.0 Hz, 1H), 5.67 (s, 1H), 4.17 (d, J=5.4 Hz, 2H), 3.92 (t, J=5.4 Hz, 1H), 2.92 (s, 12H);

MS (DCI/NH₃) m/e 460 (M+H)⁺.

WHAT IS CLAIMED IS:

1. A method of selectively modulating the glucocorticoid receptor-mediated gene expression in a mammal comprising administering an effective amount of a compound of Formula I



I,

or a pharmaceutically acceptable salt or prodrug thereof, where

 R^1 is selected from

- (1) hydrogen and
- (2) -OH;

 L^1 is selected from

- (1) a covalent bond,
- (2) -O-,
- (3) -S(O)_t- where t is an integer from 0 to 2, and
- (4) -NR⁹- where R⁹ is selected from
 - (a) hydrogen and
 - (b) alkyl of one to four carbons;

 R^2 and R^3 are independently selected from

- (1) hydrogen,
- (2) an amino-protecting group,
- (3) alkyl of one to six carbons, and
- (4) alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from
 - (a) phenyl and
 - (b) -OR¹⁰ where R¹⁰ is selected from

- 5 (i) hydrogen,
 (ii) alkyl of one to six carbons,
 (iii) a hydroxy-protecting group, and
 (iv) $-C(O)R^{11}$ where R^{11} is selected from
 alkyl of one to six carbons,
 phenyl, and
 phenyl substituted with 1, 2, or 3 substituents
 independently selected from
 $-NO_2$,
 10 alkyl of one to six carbons, and
 halogen, or

R^2 and R^3 together with the nitrogen to which they are attached form a 4 to 8
 membered ring selected from the group consisting of heterocycle;

15

R^4 and R^5 are independently selected from

- (1) hydrogen,
 (2) halogen,
 (3) $-NR^{12}R^{13}$ where R^{12} and R^{13} are independently selected from
 20 (a) hydrogen,
 (b) an amino-protecting group,
 (c) alkyl of one to six carbons, and
 (d) alkyl of one to six carbons substituted with 1, 2, or 3 substituents
 independently selected from
 25 (i) $-OR^{10}$ and
 (ii) phenyl, or
 R^{12} and R^{13} together with the nitrogen to which they are attached form a
 4 to 8 membered ring selected from the group consisting of heterocycle,
 (4) alkyl of one to six carbons, and
 30 (5) alkyl of one to six carbons substituted with 1, 2, or 3 substituents
 independently selected from
 (a) halogen,
 (b) $-OR^{10}$,
 (c) $-CN$, and

- (d) $-\text{CO}_2\text{R}^{14}$ where R^{14} is selected from
- (i) hydrogen,
 - (ii) alkyl of one to six carbons, and
 - (iii) alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from
 - phenyl and
 - phenyl substituted with 1, 2, or 3 substituents independently selected from
 - $-\text{NO}_2$,
 - alkyl of one to six carbons, and
 - halogen, and
- (e) $-\text{NR}^{15}\text{R}^{16}$ where R^{15} and R^{16} are independently selected from
- (i) hydrogen,
 - (ii) an amino-protecting group,
 - (iii) alkyl of one to six carbons, and
 - (iv) alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from
 - phenyl and
 - phenyl substituted with 1, 2, or 3 substituents independently selected from
 - $-\text{NO}_2$,
 - alkyl of one to six carbons, and
 - halogen;
- R^6 , R^7 , and R^8 are independently selected from
- (1) hydrogen,
 - (2) halogen,
 - (3) alkyl of one to six carbons,
 - (4) alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from
 - (a) halogen,
 - (b) $-\text{OR}^{10}$,
 - (c) $-\text{CN}$,
 - (d) $-\text{CO}_2\text{R}^{14}$, and

- (e) -NR¹⁵R¹⁶,
(5) perfluoroalkyl of one to six carbons,
(6) -NR¹⁷R¹⁸ where R¹⁷ and R¹⁸ are independently selected from
(a) hydrogen,
5 (b) an amino-protecting group,
(c) alkyl of one to six carbons,
(d) alkyl of one to six carbons substituted with 1, 2, or 3 substituents
independently selected from
(i) phenyl,
10 (ii) heterocycle, and
(iii) -OR¹⁰,
(e) -C(O)R¹⁹ where R¹⁹ is selected from
(i) alkyl of one to six carbons,
(ii) alkyl of one to six carbons substituted with 1, 2, or 3
15 substituents independently selected from
heterocycle,
phenyl, and
phenyl substituted with 1, 2, or 3 substituents
independently selected from
20 alkyl of one to six carbons,
halogen,
-NO₂,
-CF₃,
-CN,
25 -C(O)R²⁰ where R²⁰ is selected from
hydrogen,
alkyl of one to six carbons,
-NR²¹R²² where R²¹ and R²² are
independently selected from
30 hydrogen,
an amino-protecting group,
and
alkyl of one to six carbons,
and

-OR²³ where R²³ is selected from
alkyl of one to six carbons and
alkyl of one to six carbons substituted
with 1, 2, or 3 substituents
independently selected from the group
consisting of halogen,

5

- (iii) cycloalkyl of three to six carbons,
- (iv) cycloalkyl of three to six carbons substituted with 1, 2, or 3 substituents independently selected from

10

heterocycle,
phenyl,
phenyl substituted with 1, 2, or 3 substituents
independently selected from

15

-NO₂,
alkyl of one to six carbons, and
halogen,

halogen,

-CN,

$$-\text{CO}_2\text{R}^{14},$$

20

alkyl of one to six carbons, and
alkyl of one to six carbons substituted with 1, 2, or 3
substituents independently selected from the group
consisting of halogen,

- (v) alkenyl of two to six carbons,

25

- (vi) alkenyl of two to six carbons substituted with 1 or 2 substituents independently selected from

heterocycle,

phenyl, and

phenyl substituted with 1, 2, or 3 substituents
independently selected from

30

alkyl of one to six carbons,

halogen,

-NO₂,

 $-\text{CF}_3,$

- CN, and
-CO₂R¹⁴,
- (vii) phenyl,
(viii) phenyl substituted with 1, 2, or 3 substituents independently
5 selected from
halogen,
alkyl of one to six carbons,
-NO₂,
-CF₃,
10 -CN, and
-CO₂R¹⁴, and
- (ix) -OR¹¹, and
(f) -SO₂R²⁴ where R²⁴ is selected from
(i) alkyl of one to six carbons,
15 (ii) phenyl, and
(iii) phenyl substituted with 1, 2, or 3 substituents independently
selected from
alkyl of one to six carbons and
-NO₂,
- 20 (7) -OR²⁵ where R²⁵ is selected from
(a) perfluoroalkyl of one to six carbons,
(b) alkyl of one to six carbons,
(c) alkyl of one to six carbons substituted with 1, 2, or 3 substituents
independently selected from
25 (i) halogen and
(ii) phenyl,
(d) a hydroxy-protecting group, and
(e) -C(O)R¹⁴,
- (8) -CN,
30 (9) -C(O)R¹⁹,
(10) -CO₂R¹⁴,
(11) -C(O)R²⁰,
(12) -SO₂NR²⁶R²⁷ where R²⁶ and R²⁷ are independently selected from,
(a) alkyl of one to six carbons,

- (b) phenyl, and
- (c) phenyl substituted with 1, 2, or 3 substituents independently selected from
- (i) alkyl of one to six carbons,
- (ii) halogen, and
- (iii) $-\text{NO}_2$,
- (13) $-\text{S}(\text{O})_t\text{R}^{28}$ where t is defined previously and R^{28} is selected from
- (a) hydrogen,
- (b) perfluoroalkyl of one to six carbons,
- (c) alkyl of one to six carbons,
- (d) alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from
- (i) phenyl and
- (ii) phenyl substituted with 1, 2, or 3 substituents independently selected from
- halogen,
- alkyl of one to six carbons, and
- $-\text{NO}_2$,
- (e) phenyl, and
- (f) phenyl substituted with 1, 2, or 3 substituents independently selected from
- (i) halogen,
- (ii) alkyl of one to six carbons, and
- (iii) $-\text{NO}_2$,
- (14) $-\text{NO}_2$,
- (15) $-\text{N}=\text{CHR}^{29}$ where R^{29} is selected from
- (a) phenyl,
- (b) aryl, and
- (c) heterocycle, and
- (16)
- $$\begin{array}{l} \text{X} \\ \text{Y} \\ \text{O} \end{array}$$
- where X is selected from $-\text{CH}_2-$, $-\text{CH}_2\text{O}-$ and $-\text{O}-$, and Y is selected from $-\text{C}(\text{O})-$ and $-(\text{C}(\text{R}'')_2)_v-$, where R'' is hydrogen or alkyl of one


to four carbons, and v is an integer from 1 to 3.

2. A method of treating inflammation and immune, autoimmune, and inflammatory diseases in a mammal comprising administering an effective amount of a compound of Formula I.
3. A method of treating adrenal imbalance in a mammal comprising administering an effective amount of a compound of Formula I.
4. A method of treating cognitive and behavioral processes comprising administering an effective amount of a compound of Formula I.
5. A method according to Claim 4 where the cognitive or behavioral process is selected from cognitive performance, memory and learning enhancement, depression, addiction, mood disorders, chronic fatigue syndrome, schizophrenia, stroke, sleep disorders, and anxiety.
6. The method of claim 1 where the compound of Formula I is selected from the group consisting of
 - 4',4'' bis(dimethylamino)-2-chloro-5-nitrotriphenylmethane,
 - 4',4'' bis(dimethylamino)-4-chloro-3-nitrotriphenylmethane,
 - 5 4',4'' bis(dimethylamino)-5-acetamido-2-chlorotriphenylmethane,
 - 4',4'' bis(dimethylamino)-4-nitrotriphenylmethane,
 - 4',4'' bis(dimethylamino)-4-chlorotriphenylmethane,
 - 4',4'' bis(dimethylamino)-3-chlorotriphenylmethane,
 - 4',4'' bis(dimethylamino)-2-chlorotriphenylmethane,
 - 10 4',4'' bis(dimethylamino)-2-methoxytriphenylmethane,
 - 4',4'' bis(dimethylamino)-3-nitrotriphenylmethane,
 - 4',4'' bis(dimethylamino)-2-trifluoromethyltriphenylmethane,
 - 4',4'' bis(dimethylamino)triphenylmethane,
 - 4',4'' bis(dimethylamino)-2-chloro-6-nitrotriphenylmethane,
 - 15 4',4'' bis(N-piperidiny)-2-chloro-5-nitrotriphenylmethane,
 - 4',4'' bis(dimethylamino)-2-bromotriphenylmethane,
 - 4',4'' bis(dimethylamino)-2-methyltriphenylmethane,

- 4',4'' bis(dimethylamino)-2,3,5-trichlorotriphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-5-trifluoromethyltriphenylmethane,
20 4',4'' bis(dimethylamino)-2,4-dichlorotriphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-4,5-methylenedioxytriphenylmethane,
4',4'' bis(dimethylamino)-2,6-dichlorotriphenylmethane,
4',4'' bis(dimethylamino)-2,3-dichlorotriphenylmethane,
4',4'' bis(N-morpholinyl)-2-chloro-5-nitrotriphenylmethane,
25 4',4'' bis(methylamino)-2-chloro-5-nitrotriphenylmethane,
4',4'' bis(dimethylamino)-2,5-dichlorotriphenylmethane,
4',4'' bis(dimethylamino)-2-fluorotriphenylmethane,
4',4'' bis(dimethylamino)-2-iodotriphenylmethane,
4',4'' bis(methylamino)-2,5-dichlorotriphenylmethane,
30 4',4'' bis(N-morpholinyl)-2,3,5-trichlorotriphenylmethane,
4',4'' bis(N-pyrrolidinyl)-2-chloro-5-nitrotriphenylmethane,
4',4'' bis(di-n-butylamino)-2-chloro-5-nitrotriphenylmethane,
4',4'' bis(N-(2-acetoxyethyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane,
4',4'' bis(N-(2-hydroxyethyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane,
35 4',4'' bis(dimethylamino)-2-chloro-5-iodotriphenylmethane,
4',4'' bis(dimethylamino)-5-bromo-2-chlorotriphenylmethane,
4',4'' bis(N-(t-butoxycarbonyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane,
4',4'' bis(N-benzylamino)-2-chloro-5-nitrotriphenylmethane,
4',4'' bis(N-benzylamino)-2,5-dichlorotriphenylmethane,
40 4',4'' bis(dimethylamino)-4-methoxytriphenylmethane,
4'-dimethylamino-4''-methylamino-2-chloro-5-nitrotriphenylmethane,
4'-dimethylamino-4''-(N-morpholinyl)-2-chloro-5-nitrotriphenylmethane,
4'-dimethylamino-4''-(N-pyrrolidinyl)-2-chloro-5-nitrotriphenylmethane,
4'-dimethylamino-4''-(N-methyl-N-(2-hydroxyethyl)amino)-2-chloro-5-
45 nitrotriphenylmethane,
4'-dimethylamino-4''-(di-n-butylamino)-2-chloro-5-nitrotriphenylmethane,
3'-methyl-4',4''-bis(dimethylamino)-2-chloro-5-nitrotriphenylmethane,
4'-dimethylamino-4''-(N-piperidinyl)-2-chloro-5-nitrotriphenylmethane,
4'-methylamino-4''-(t-butoxycarbonylamino)-2-chloro-5-nitrotriphenylmethane,
50 4'-methylamino-4''-(N-(2-acetoxyethyl)-N-methylamino)-2-chloro-5-
nitrotriphenylmethane,

4'-dimethylamino-4''-(N-methyl-N-(2-benzoyloxyethyl)amino)-2-chloro-5-nitrotriphenylmethane,
4'-dimethylamino-4''-(N-methyl-N-(2-acetoxyethyl)amino)-2-chloro-5-nitrotriphenylmethane,
55 phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] ether,
4-chlorophenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] ether,
phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] thioether,
phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] amine,
60 4'-dimethylamino-2-chloro-5-nitrotriphenylmethane,
4'-dimethylamino-2-chlorotriphenylmethanol,
4'-dimethylamino-2-chlorotriphenylmethane,
4',4'' bis(dimethylamino)-5-amino-2-chlorotriphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-5-(4-nitrobenzamido)triphenylmethane,
65 4',4'' bis(dimethylamino)-2-chloro-5-(4-nitrocinnamido)triphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-5-(cyclopropylcarbamido)triphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-5-(dimethylsulphonimido)triphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-5-(methoxycarbonylamino)triphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-5-(2-furanylmethylimino)triphenylmethane, and
70 4',4'' bis(dimethylamino)-2-chloro-5-(2-furanylmethylamino)triphenylmethane.

7. A pharmaceutical composition comprising a therapeutically-effective amount of a compound of Formula I in combination with a pharmaceutically acceptable carrier.
8. A compound of Formula II



II,

5 or a pharmaceutically acceptable salt or prodrug thereof, where

R¹ is selected from

- (1) hydrogen and
- (2) -OH;

10

L¹ is selected from

- (1) a covalent bond,
- (2) -O-,
- (3) -S(O)_t- where t is an integer from 0 to 2, and
- (4) -NR⁹- where R⁹ is selected from
 - (a) hydrogen and
 - (b) alkyl of one to four carbons;

15

R² and **R³** are independently selected from

- (1) hydrogen,
- (2) an amino-protecting group,
- (3) alkyl of one to six carbons, and
- (4) alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from
 - (a) phenyl and
 - (b) -OR¹⁰ where R¹⁰ is selected from

25

- 30 (i) hydrogen,
 (ii) alkyl of one to six carbons,
 (iii) a hydroxy-protecting group, and
 (iv) $-C(O)R^{11}$ where R^{11} is selected from
 alkyl of one to six carbons,
 phenyl, and
 phenyl substituted with 1, 2, or 3 substituents
 independently selected from
 35 $-NO_2$,
 alkyl of one to six carbons,
 halogen, or

40 R^2 and R^3 together with the nitrogen to which they are attached form a 4 to 8
 membered ring selected from the group consisting of heterocycle;

R^4 and R^5 are independently selected from

- (1) hydrogen,
 (2) halogen,
 45 (3) $-NR^{12}R^{13}$ where R^{12} and R^{13} are independently selected from
 (a) hydrogen,
 (b) an amino-protecting group,
 (c) alkyl of one to six carbons, and
 (d) alkyl of one to six carbons substituted with 1, 2, or 3 substituents
 50 independently selected from
 (i) $-OR^{10}$ and
 (ii) phenyl, or
 R^{12} and R^{13} together with the nitrogen to which they are attached form a
 4 to 8 membered ring selected from the group consisting of heterocycle,
 55 (4) alkyl of one to six carbons, and
 (5) alkyl of one to six carbons substituted with 1, 2, or 3 substituents
 independently selected from
 (a) halogen,
 (b) $-OR^{10}$,
 60 (c) $-CN$,

- (d) $-\text{CO}_2\text{R}^{14}$ where R^{14} is selected from
- (i) hydrogen,
 - (ii) alkyl of one to six carbons, and
 - (iii) alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from
phenyl and
phenyl substituted with 1, 2, or 3 substituents
independently selected from
 $-\text{NO}_2$,
alkyl of one to six carbons, and
halogen, and
- (e) $-\text{NR}^{15}\text{R}^{16}$ where R^{15} and R^{16} are independently selected from
- (i) hydrogen,
 - (ii) an amino-protecting group,
 - (iii) alkyl of one to six carbons, and
 - (iv) alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from
phenyl and
phenyl substituted with 1, 2, or 3 substituents
independently selected from
 $-\text{NO}_2$,
alkyl of one to six carbons, and
halogen;
- R^6 is selected from the group consisting of halogen;
- R^7 is selected from
- (1) $-\text{NO}_2$,
 - (2) $-\text{CF}_3$,
 - (3) $-\text{NR}^{17}\text{R}^{18}$ where R^{17} and R^{18} are independently selected from
 - (a) hydrogen,
 - (b) an amino-protecting group,
 - (c) alkyl of one to six carbons,
 - (d) alkyl of one to six carbons substituted with 1, 2, or 3 substituents

- 95 independently selected from
- (i) phenyl,
 - (ii) heterocycle, and
 - (iii) $-OR^{10}$,
- (e) $-C(O)R^{19}$ where R^{19} is selected from
- 100 (i) alkyl of one to six carbons,
- (ii) alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from
- heterocycle,
- phenyl, and
- 105 phenyl substituted with 1, 2, or 3 substituents independently selected from
- alkyl of one to six carbons,
- halogen,
- $-NO_2$,
- 110 $-CF_3$,
- $-CN$,
- $-C(O)R^{20}$ where R^{20} is selected from
- hydrogen,
- alkyl of one to six carbons, and
- 115 $-NR^{21}R^{22}$ where R^{21} and R^{22} are independently selected from
- hydrogen,
- an amino-protecting group,
- and
- 120 alkyl of one to six carbons,
- and
- $-OR^{23}$ where R^{23} is selected from
- alkyl of one to six carbons and
- alkyl of one to six carbons substituted with 1, 2, or 3 substituents independently selected from the group consisting of halogen,
- 125 (iii) cycloalkyl of three to six carbons,

- 130 (iv) cycloalkyl of three to six carbons substituted with 1, 2, or 3 substituents independently selected from
heterocycle,
phenyl,
phenyl substituted with 1, 2, or 3 substituents
independently selected from
- 135 -NO₂,
alkyl of one to six carbons, and
halogen,
halogen,
-CN,
140 -CO₂R¹⁴,
alkyl of one to six carbons, and
alkyl of one to six carbons substituted with 1, 2, or 3
substituents independently selected from the group
consisting of halogen,
- 145 (v) alkenyl of two to six carbons,
(vi) alkenyl of two to six carbons substituted with 1 or 2
substituents independently selected from
heterocycle,
phenyl, and
150 phenyl substituted with 1, 2, or 3 substituents
independently selected from
alkyl of one to six carbons,
halogen,
-NO₂,
155 -CF₃,
-CN, and
-CO₂R¹⁴,
- (vii) phenyl,
(viii) phenyl substituted with 1, 2, or 3 substituents independently
160 selected from
halogen,
alkyl of one to six carbons,

-NO₂,
-CF₃,
-CN, and
-CO₂R¹⁴, and

165

(f) -OR¹¹, and
 -SO₂R²⁴ where R²⁴ is selected from
 (i) alkyl of one to six carbons,
 (ii) phenyl, and
 (iii) phenyl substituted with 1, 2, or 3 substituents independently selected from
 alkyl of one to six carbons and
 -NO₂, and

170

175 (4) -N=CHR²⁹ where R²⁹ is selected from

- (a) phenyl,
- (b) aryl, and
- (c) heterocycle;

180 **R⁸** is selected from

(1) hydrogen and

(2) halogen; or

R⁶ and R⁸ or R⁷ and R⁸ taken together can form

185

$\begin{matrix} \text{X} \\ | \\ \text{---} \end{matrix}$
 $\begin{matrix} \text{Y} \\ | \\ \text{---} \end{matrix}$
where X is selected from -CH₂-, -CH₂O- and -O-, and Y is selected from -C(O)- and -(C(R''))_v -, where R'' is hydrogen or alkyl of one to four carbons, and v is an integer from 1 to 3.

9. A compound according to claim 8 where R^6 is -Cl, R^7 is -NO₂, and R^8 is hydrogen.

10. A compound according to claim 9 selected from the group consisting of
4',4" bis(dimethylamino)-2-chloro-5-nitrotriphenylmethane,
4',4" bis(N-piperidiny1)-2-chloro-5-nitrotriphenylmethane,
4',4" bis(N-morpholiny1)-2-chloro-5-nitrotriphenylmethane,

- 4',4'' bis(methylamino)-2-chloro-5-nitrotriphenylmethane,
 4',4'' bis(N-pyrrolidinyl)-2-chloro-5-nitrotriphenylmethane,
 4',4'' bis(di-n-butylamino)-2-chloro-5-nitrotriphenylmethane,
 4',4'' bis(N-(2-acetoxyethyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane,
 4',4'' bis(N-(2-hydroxyethyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane,
 4',4'' bis(N-(t-butoxycarbonyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane,
 4',4'' bis(N-benzylamino)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-methylamino-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-morpholinyl)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-pyrrolidinyl)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-methyl-N-(2-hydroxyethyl)amino)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(di-n-butylamino)-2-chloro-5-nitrotriphenylmethane,
 3'-methyl-4',4''-bis(dimethylamino)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-piperidinyl)-2-chloro-5-nitrotriphenylmethane,
 4'-methylamino-4''-(t-butoxycarbonylamino)-2-chloro-5-nitrotriphenylmethane,
 4'-methylamino-4''-(N-(2-acetoxyethyl)-N-methylamino)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-methyl-N-(2-benzoyloxyethyl)amino)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-methyl-N-(2-acetoxyethyl)amino)-2-chloro-5-nitrotriphenylmethane,
 phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] ether,
 4-chlorophenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] ether,
 phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] thioether,
 phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] amine, and
 4'-dimethylamino-2-chloro-5-nitrotriphenylmethane.
11. A compound according to claim 8 where R^6 is -Cl, R^7 is -CF₃, and R^8 is hydrogen.
 12. A compound according to claim 11 that is
 4',4'' bis(dimethylamino)-2-chloro-5-trifluoromethyltriphenylmethane.
 13. A compound according to claim 8 where R^6 is -Cl, R^7 is -NR¹⁷R¹⁸, and R^8 is hydrogen.

14. A compound according to claim 13 selected from the group consisting of
 4',4'' bis(dimethylamino)-5-acetamido-2-chlorotriphenylmethane,
 4',4'' bis(dimethylamino)-5-amino-2-chlorotriphenylmethane,
 4',4'' bis(dimethylamino)-2-chloro-5-(4-nitrobenzamido)triphenylmethane,
 5 4',4'' bis(dimethylamino)-2-chloro-5-(4-nitrocinnamido)triphenylmethane,
 4',4'' bis(dimethylamino)-2-chloro-5-(cyclopropylcarbamido)triphenylmethane,
 4',4'' bis(dimethylamino)-2-chloro-5-(dimethylsulphonimido)triphenylmethane,
 4',4',4'' bis(dimethylamino)-2-chloro-5-(methoxycarbonylamino)triphenylmethane,
 10 4',4'' bis(dimethylamino)-2-chloro-5-(2-furanylmethylamino)triphenylmethane.
15. A compound according to claim 8 where R^6 is -Cl, R^7 is -N=CHR²⁹, and R^8 is hydrogen.
16. A compound according to claim 15 that is
 4',4'' bis(dimethylamino)-2-chloro-5-(2-furanylmethylimino)triphenylmethane.
17. A compound according to claim 8 where R^6 is -Cl and R^7 and R^8 taken together form
- $$\begin{array}{c} \xi - X \\ \xi - O - Y \end{array}$$
- where X is -O- and Y is -(C(R'')₂)_v - where R'' is hydrogen and v is 1.
18. A compound according to claim 17 that is
 4',4'' bis(dimethylamino)-2-chloro-4,5-methylenedioxytriphenylmethane.

19. A compound selected from the group consisting of
- 4',4'' bis(dimethylamino)-2-chloro-5-nitrotriphenylmethane,
 4',4'' bis(dimethylamino)-5-acetamido-2-chlorotriphenylmethane,
 4',4'' bis(N-piperidinyl)-2-chloro-5-nitrotriphenylmethane,
 5 4',4'' bis(dimethylamino)-2-chloro-5-trifluoromethyltriphenylmethane,
 4',4'' bis(dimethylamino)-2-chloro-4,5-methylenedioxytriphenylmethane,
 4',4'' bis(N-morpholinyl)-2-chloro-5-nitrotriphenylmethane,
 4',4'' bis(methylamino)-2-chloro-5-nitrotriphenylmethane,
 4',4'' bis(N-pyrrolidinyl)-2-chloro-5-nitrotriphenylmethane,
 10 4',4'' bis(di-n-butylamino)-2-chloro-5-nitrotriphenylmethane,
 4',4'' bis(N-(2-acetoxyethyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane,
 4',4'' bis(N-(2-hydroxyethyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane,
 4',4'' bis(N-(t-butoxycarbonyl) N-methylamino)-2-chloro-5-nitrotriphenylmethane,
 4',4'' bis(N-benzylamino)-2-chloro-5-nitrotriphenylmethane,
 15 4'-dimethylamino-4''-methylamino-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-morpholinyl)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-pyrrolidinyl)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-methyl-N-(2-hydroxyethyl)amino)-2-chloro-5-nitrotriphenylmethane,
 20 4'-dimethylamino-4''-(di-n-butylamino)-2-chloro-5-nitrotriphenylmethane,
 3'-methyl-4',4''-bis(dimethylamino)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-piperidinyl)-2-chloro-5-nitrotriphenylmethane,
 4'-methylamino-4''-(t-butoxycarbonylamino)-2-chloro-5-nitrotriphenylmethane,
 4'-methylamino-4''-(N-(2-acetoxyethyl)-N-methylamino)-2-chloro-5-nitrotriphenylmethane,
 25 4'-dimethylamino-4''-(N-methyl-N-(2-benzoyloxyethyl)amino)-2-chloro-5-nitrotriphenylmethane,
 4'-dimethylamino-4''-(N-methyl-N-(2-acetoxyethyl)amino)-2-chloro-5-nitrotriphenylmethane,
 30 phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] ether,
 4-chlorophenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] ether,
 phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] thioether,
 phenyl-[(4'-dimethylaminophenyl)-(2-chloro-5-nitrophenyl)methyl] amine,
 4'-dimethylamino-2-chloro-5-nitrotriphenylmethane,

- 35 4',4'' bis(dimethylamino)-5-amino-2-chlorotriphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-5-(4-nitrobenzamido)triphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-5-(4-nitrocinnamido)triphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-5-(cyclopropylcarbarnido)triphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-5-(dimethylsulphonimido)triphenylmethane,
40 4'4',4'' bis(dimethylamino)-2-chloro-5-(methoxycarbonylamino)triphenylmethane,
4',4'' bis(dimethylamino)-2-chloro-5-(2-furanylmethylimino)triphenylmethane, and
4',4'' bis(dimethylamino)-2-chloro-5-(2-furanylmethylamino)triphenylmethane.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/136, 31/137, 31/167, 31/4545, <i>(continued on the following page)</i>	A3	(11) International Publication Number: WO 00/06137 (43) International Publication Date: 10 February 2000 (10.02.00)
(21) International Application Number: PCT/US99/17267 (22) International Filing Date: 30 July 1999 (30.07.99) (30) Priority Data: 09/126,185 30 July 1998 (30.07.98) US (71) Applicant: ABBOTT LABORATORIES [US/US]; CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-6050 (US). (72) Inventors: COGHLAN, Michael, J.; 307 Fairfax Lane, Grayslake, IL 60030 (US). LULY, Jay, R.; 24 Damien Road, Wellesley, MA 02181 (US). SCHKERYANTZ, Jeffrey, M.; 526 English Lane, Winthrop Harbor, IL 60096 (US). WANG, Alan, X.; 1012 Talbot Lane, Grayslake, IL 60030 (US). (74) Agents: MILLER, Robert, A. et al.; Abbott Laboratories, CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US).		(81) Designated States: CA, JP, MX, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report</i> <i>Before the expiration of the time limit for amending the claims</i> <i>and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 27 April 2000 (27.04.00)
(54) Title: GLUCOCORTICOID-SELECTIVE ANTI-INFLAMMATORY AGENTS <div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div> (57) Abstract Compounds having Formula (I) are useful for modulating the glucocorticoid receptor in a mammal. Also disclosed are pharmaceutical compositions comprising compounds of Formula (I) and methods of treating immune, autoimmune, inflammatory, adrenal imbalance, cognitive and behavioral diseases in a mammal.		

A61K 31/535, 31/5375, 31/5377, 31/4025, 31/222, 31/27, 31/40, 31/45, 31/24, 31/341, C07D 295/06, C07C 219/06, 211/52, 215/16, 271/28, 219/10, 217/76, 323/29, 233/43, 233/80, 233/40, 233/62, C07D 307/52, C07C 317/58

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/17267

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/136 A61K31/137 A61K31/167 A61K31/4545 A61K31/535
A61K31/5375 A61K31/5377 A61K31/4025 A61K31/222 A61K31/27
A61K31/40 A61K31/45 A61K31/24 A61K31/341 C07D295/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07D C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 99 32101 A (ALCON LAB INC ; CLARK ABBOT F (US)) 1 July 1999 (1999-07-01) claims 1,2,5	1,7
X	WO 97 34589 A (ION PHARMACEUTICALS INC ; HARVARD COLLEGE (US); CHILDRENS MEDICAL C) 25 September 1997 (1997-09-25) * See compounds RN= 467-62-9 * page 87, line 25 -page 88, line 15 -/--	7

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 February 2000

Date of mailing of the international search report

29/02/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Veronese, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/17267

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07C219/06 C07C211/52 C07C215/16 C07C271/28 C07C219/10 C07C217/76 C07C323/29 C07C233/43 C07C233/80 C07C233/40 C07C233/62 C07D307/52 C07C317/58		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols)		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US MITSUBISHI PAPER MILLS, LTD., JAPAN: "Electrophotographic photosensitive material" retrieved from STN Database accession no. 102:70133 XP002130786 * See compound RN=94497-79-7 * abstract & JP 59 036255 A (MITSUBISHI PAPER MILLS, LTD., JAPAN) 28 February 1984 (1984-02-28)	8
X	DE 25 03 309 A (CIBA GEIGY AG) 7 August 1975 (1975-08-07) * See compounds RN= 57752-10-0 and 6310-52-7 (N. 62 and N. 33) * --- -/--	8
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
<div style="display: flex;"> <div style="flex: 1;"> <p>* Special categories of cited documents</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center;">17 February 2000</div>		Date of mailing of the international search report
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center;">Veronese, A</div>

INTERNATIONAL SEARCH REPORT

Inter. Patent Application No.

PCT/US 99/17267

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 395 093 A (KYOWA HAKKO KOGYO KK) 31 October 1990 (1990-10-31) cited in the application the whole document ---	1-18
A	US 4 152 341 A (JONES GERAINT ET AL) 1 May 1979 (1979-05-01) the whole document ---	1-19
A	LOOSE D S ET AL: "Ketoconazole Binds to Glucocorticoid Receptors and Exhibits Glucocorticoid Antagonist Activity in Cultured Cells" JOURNAL OF CLINICAL INVESTIGATION,US,NEW YORK, NY, vol. 72, 1 July 1983 (1983-07-01), pages 404-408, XP002095830 ISSN: 0021-9738 -----	1-19

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/17267

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 1-6
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/17267

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9932101 A	01-07-1999	AU 1714399 A	12-07-1999
WO 9734589 A	25-09-1997	AU 2538897 A CA 2250092 A EP 0918514 A	10-10-1997 25-09-1997 02-06-1999
JP 59036255 A	28-02-1984	NONE	
DE 2503309 A	07-08-1975	GB 1469481 A AT 334406 B AT 71775 A BE 825040 A CH 594509 A ES 434322 A FI 750166 A FR 2259701 A IT 1029406 B JP 50108014 A NL 7501191 A SE 7500447 A US 3995088 A ZA 7500670 A	06-04-1977 10-01-1976 15-05-1976 31-07-1975 13-01-1978 16-12-1976 02-08-1975 29-08-1975 10-03-1979 26-08-1975 05-08-1975 04-08-1975 30-11-1976 28-01-1976
EP 0395093 A	31-10-1990	AT 93838 T CA 2015473 A,C DE 69002996 D JP 2749951 B JP 3215461 A US 5413997 A US 5112867 A	15-09-1993 28-10-1990 07-10-1993 13-05-1998 20-09-1991 09-05-1995 12-05-1992
US 4152341 A	01-05-1979	GB 1560880 A AT 767577 A AT 346835 B AT 971676 A AU 512090 B AU 2060576 A BE 849872 A CA 1086743 A CH 627156 A CH 624378 A CS 191326 B CY 1147 A DD 128671 A DE 2658840 A DK 586576 A,B, ES 454657 A FI 763735 A FR 2336921 A HK 61881 A HU 176679 B IE 43994 B IL 51065 A IN 145508 A JP 1350116 C JP 52083356 A JP 61015061 B KE 3169 A	13-02-1980 15-11-1978 27-11-1978 15-04-1978 25-09-1980 22-06-1978 24-06-1977 30-09-1980 31-12-1981 31-07-1981 29-06-1979 10-09-1982 30-11-1977 07-07-1977 30-06-1977 16-03-1978 30-06-1977 29-07-1977 18-12-1981 28-04-1981 15-07-1981 31-12-1980 28-10-1978 28-11-1986 12-07-1977 22-04-1986 06-11-1981

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 99/17267

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4152341 A		LU 76483 A	18-01-1978
		MX 4655 E	19-07-1982
		NL 7614015 A	01-07-1977
		NO 764371 A,B,	30-06-1977
		NZ 182807 A	03-04-1978
		PH 13545 A	26-06-1980
		SE 428468 B	04-07-1983
		SE 7614304 A	30-06-1977
		US RE31467 E	20-12-1983
		US 4113879 A	12-09-1978
		ZA 7607215 A	30-11-1977

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.